


# Basic ideas of hypothesis testing



All examples are fictitious. All data are simulated and the graphics were created with the statistical program package R.

The materials are protected by copyright and are only provided for personal use for studies at TU Vienna. Further use is not permitted. In particular, it is not permitted to distribute the materials or make them publicly available (e.g. in social networks, on learning platforms, etc.).

Sämtliche Beispiele sind frei erfunden. Alle Daten sind simuliert und die Grafiken wurden mit statistischen Programmpaket R erstellt.


Die Materialien sind urheberrechtlich geschützt und dürfen ausschließlich für den Eigengebrauch im Rahmen des Studiums an der TU Wien genutzt werden. Eine weitere Nutzung ist nicht gestattet. Insbesondere ist es nicht gestattet, die Materialien zu verbreiten oder öffentlich zugänglich zu machen (etwa im Rahmen sozialer Netzwerke, Lernplattformen etc.).

# Reminder

- How is the **mean** distributed under normal distribution?


# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$




# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$




# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed




# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )



# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$






# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$


(n=16)



# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$


(n=16)



# Reminder


- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$

(n=16)




# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$




# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$




# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$



# Reminder


- How is the **mean** distributed under normal distribution?
  - Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
  - For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
    - $\bar{X}$  is also normally distributed
    - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
    - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )
- Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$



# Reminder

- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )


Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$





# Reminder


- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$



# Reminder


- How is the **mean** distributed under normal distribution?
- Let  $X_1, \dots, X_n$  be independent and identically distributed (i.i.d.) random variables and  $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds  $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$ 
  - $\bar{X}$  is also normally distributed
  - $\bar{X}$  has expectation  $\mu_{\bar{X}} = \mu$  (equal to the expectation of  $X_i$ )
  - $\bar{X}$  has standard deviation  $\sigma_{\bar{X}} = \sigma / \sqrt{n}$  (decrease of factor  $1/\sqrt{n}$ )Interpretation: the typical deviation of the **mean** from its expectation is  $\sigma / \sqrt{n}$

- **Standardization:**  $\frac{\bar{X} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$



# Overview


We differentiate:  
Probability theory  
(Stochastics)  
=  
Theory of randomness



# Overview

We differentiate:  
Probability theory  
(Stochastics)  
=  
Theory of randomness

and  
Statistics  
=  
Description of data →



# Overview

We differentiate:

Probability theory  
(Stochastics)

=



Theory of randomness

and

Statistics

=

Description of data →  
(using stochastic **models**)




# Overview

We differentiate:

Probability theory  
(Stochastics)

=

Theory of randomness




and

Statistics

=

Description of data →  
(using stochastic **models**)



---

Previous lecture: Short excursion to descriptive Statistics

How do data look like? How can they be summarized?

# Overview

We differentiate:

Probability theory  
(Stochastics)

=



Theory of randomness

and

Statistics

=

Description of data →  
(using stochastic **models**)



---

Previous lecture: Short excursion to descriptive Statistics

How do data look like? How can they be summarized?


Today: inferentiell Statistics (Modelling)

How did the data occur?

# Basic problem

Reminder:

- Runtimes of an algorithm implemented by  $n_A = 121$  students






# Basic problem

Reminder:


- Runtimes of an algorithm implemented by  $n_A = 121$  students
- Additionally implemented from  $n_B = 16$  students that took a certain programming course



# Basic problem

Reminder:

- Runtimes of an algorithm implemented by  $n_A = 121$  students
- Additionally implemented from  $n_B = 16$  students that took a certain programming course




- Distributions shifted against each other → the course seems to have a positive effect...

# Basic problem


Reminder:

- Runtimes of an algorithm implemented by  $n_A = 121$  students
- Additionally implemented from  $n_B = 16$  students that took a certain programming course




- Distributions shifted against each other → the course seems to have a positive effect...
- In the following consider only group *B* (*one-sample situation*)

# Basic problem




- Distributions shifted against each other → the course seems to have a positive effect...

# Basic problem




- Distributions shifted against each other → the course seems to have a positive effect...
- ...the lecturer is happy!

# Basic problem



- Distributions shifted against each other → the course seems to have a positive effect...
- ...the lecturer is happy!
- A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!"

# Basic problem




- Distributions shifted against each other → the course seems to have a positive effect...
- ...the lecturer is happy!
- A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!" And further he claims:  
"The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."



huge population  
(unknown)


- Problem: Assertion about a huge unknown *population*



# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."




- Problem: Assertion about a huge unknown population
- However, a subset known: the sample  $x_1, \dots, x_n$

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."




- Problem: Assertion about a huge unknown population
- However, a subset known: the *sample*  $x_1, \dots, x_n$
- Main questions:

How 'compatible' are the **data** with the **assertion**?

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."



- Problem: Assertion about a huge unknown population
- However, a subset known: the *sample*  $x_1, \dots, x_n$
- Main questions:

How 'compatible' are the **data** with the **assertion**?


- To do:  
Quantification of the 'discrepancy' of the data and the assertion

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

**runtimes**




- Main questions:
  - How 'compatible' are the **data** with the **assertion**?
- To do:
  - 'Quantification of the discrepancy' between **data** and **assertion**

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

**runtimes**



- Main questions:

How 'compatible' are the **data** with the **assertion**?

- To do:


- 'Quantification of the discrepancy' between **data** and **assertion**
- For example through the difference  $d = \bar{x} - \mu_0$

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

**runtimes**



- Main questions:

How 'compatible' are the **data** with the **assertion**?

- To do:


- 'Quantification of the discrepancy' between **data** and **assertion**
- For example through the difference  $d = \bar{x} - \mu_0$
- Line of thought:  $|d|$  large  $\leftrightarrow$  data hardly compatible with the assertion

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

**runtimes**



- Main questions:

How 'compatible' are the **data** with the **assertion**?

- To do:


- 'Quantification of the discrepancy' between **data** and **assertion**
- For example through the difference  $d = \bar{x} - \mu_0$
- Line of thought:  $|d|$  large  $\leftrightarrow$  data hardly compatible with the assertion
- Question: What does 'large' mean?

# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

**runtimes**



- Main questions:

How 'compatible' are the **data** with the **assertion**?

- To do:

- 'Quantification of the discrepancy' between **data** and **assertion**
- For example through the difference  $d = \bar{x} - \mu_0$
- Line of thought:  $|d|$  large  $\leftrightarrow$  data hardly compatible with the assertion
- Question: What does 'large' mean? So far we do not have a notion for 'size'!




# Basic problem

Assertion:

"If all participants that have ever taken the course, had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."

**runtimes**



- Main questions:

How 'compatible' are the **data** with the **assertion**?


- To do:

- 'Quantification of the discrepancy' between **data** and **assertion**
- For example through the difference  $d = \bar{x} - \mu_0$
- Line of thought:  $|d|$  large  $\leftrightarrow$  data hardly compatible with the assertion
- Question: What does 'large' mean? So far we do not have a notion for 'size'!  
→ need the concept of the *statistical model*!

# Statistical model

- Main questions: How 'compatible' are the **data** with the assertion?
- Idea of the statistical model:


Interpret the data  $x_1, \dots, x_n$  as realizations  
of random variables  $X_1, \dots, X_n$



# Statistical model

- Main questions: How 'compatible' are the **data** with the assertion?
- Idea of the statistical model:

Interpret the data  $x_1, \dots, x_n$  as realizations  
of random variables  $X_1, \dots, X_n$




- Upside: Notion of 'size' through probability statements

# Statistical model

- Main questions: How 'compatible' are the **data** with the assertion?
- Idea of the statistical model:

Interpret the data  $x_1, \dots, x_n$  as realizations  
of random variables  $X_1, \dots, X_n$




- Upside: Notion of 'size' through probability statements
  - Observe: almost all **data** lie in the left tail of the **blue distribution**

# Statistical model

- Main questions: How 'compatible' are the **data** with the assertion?
- Idea of the statistical model:

Interpret the data  $x_1, \dots, x_n$  as realizations  
of random variables  $X_1, \dots, X_n$




- Upside: Notion of 'size' through probability statements
  - Observe: almost all **data** lie in the left tail of the **blue distribution**
  - Interpretation: Assume that the data indeed derived from independent drawings of the blue distribution, then something unlikely has happened. ( $\leftrightarrow$  incompatibility of data and assertion)

# Statistical model

- Main questions: How 'compatible' are the **data** with the assertion?
- Idea of the statistical model:

Interpret the data  $x_1, \dots, x_n$  as realizations  
of random variables  $X_1, \dots, X_n$




- Upside: Notion of 'size' through probability statements
  - Observe: almost all **data** lie in the left tail of the **blue distribution**
  - Interpretation: Assume that the data indeed derived from independent drawings of the blue distribution, then something unlikely has happened. ( $\leftrightarrow$  incompatibility of data and assertion)
  - Gain: In the context of the model, we can quantify the discrepancy via probability statements ('notion of size')

# Statistical model

- Main questions: How 'compatible' are the **data** with the assertion?
- Idea of the statistical model:

Interpret the data  $x_1, \dots, x_n$  as realizations  
of random variables  $X_1, \dots, X_n$



- Upside: Notion of 'size' through probability statements
  - Observe: almost all **data** lie in the left tail of the **blue distribution**
  - Interpretation: Assume that the data indeed derived from independent drawings of the blue distribution, then something unlikely has happened. ( $\leftrightarrow$  incompatibility of data and assertion)
  - Gain: In the context of the model, we can quantify the discrepancy via probability statements ('notion of size')
  - $\rightarrow$  *Hypothesis test*: the procedure is as follows...

# Hypothesis test, exemplary: z-Tests

- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )




# Hypothesis test, exemplary: z-Tests

- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )
- *null hypothesis*:  $H_0 : \mu = 30$   
(Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

# Hypothesis test, exemplary: z-Tests

- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )
- *null hypothesis*:  $H_0 : \mu = 30$   
(Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )
- *test statistic*: for the evaluation of the data (measures discrepancy). Here z-statistic

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$



# Hypothesis test, exemplary: z-Tests


- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )
- *null hypothesis*:  $H_0 : \mu = 30$   
(Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )
- *test statistic*: for the evaluation of the data (measures discrepancy). Here z-statistic

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

- Theoretical distribution of the test statistic if  $H_0$  is true

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$

Distribution of Z under  $H_0 : \mu = 30$



# Hypothesis test, exemplary: z-Tests


- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )
- *null hypothesis*:  $H_0 : \mu = 30$   
(Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )
- *test statistic*: for the evaluation of the data (measures discrepancy). Here z-statistic

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

Distribution of Z under  $H_0 : \mu = 30$

- Theoretical distribution of the test statistic if  $H_0$  is true

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$



- *p-value*: quantifies the discrepancy (judge  $z$  according to the distribution of  $Z$ )

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if  $H_0$  holds true

# Hypothesis test, exemplary: z-Tests

- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )
- *null hypothesis*:  $H_0 : \mu = 30$   
(Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )
- *test statistic*: for the evaluation of the data (measures discrepancy). Here z-statistic

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

Distribution of Z under  $H_0 : \mu = 30$

- Theoretical distribution of the test statistic if  $H_0$  is true

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$



- *p-value*: quantifies the discrepancy (judge  $z$  according to the distribution of  $Z$ )

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if  $H_0$  holds true

- *Decision*: Reject the null hypothesis (reason:  $p$  is small)  
We say: the observed discrepancy was *significant* ( $p < 10^{-6}$ )

# Hypothesis test, exemplary: z-Tests


- *model assumption*:  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  (here  $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )
- *null hypothesis*:  $H_0 : \mu = 30$   
(Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )
- *test statistic*: for the evaluation of the data (measures discrepancy). Here z-statistic

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

Distribution of Z under  $H_0 : \mu = 30$

- Theoretical distribution of the test statistic if  $H_0$  is true

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$



- *p-value*: quantifies the discrepancy (judge  $z$  according to the distribution of  $Z$ )

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$


Probability to make an observation which is *at least as extreme* as in the data, if  $H_0$  holds true

- *Decision*: Reject the null hypothesis (reason:  $p$  is small)  
We say: the observed discrepancy was *significant* ( $p < 10^{-6}$ )
- *Interpretation*: the  $p$ -value is negligible. If the null hypothesis holds true, then something unlikely has happened. In that sense, the data are barely compatible with the null hypothesis.

# Remark - Model assumptions

*Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  ( $n = 16$ )  
(The data are realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )

**runtimes**




- Why are the assumptions reasonable?

# Remark - Model assumptions

*Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  ( $n = 16$ )  
(The data are realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )

**runtimes**




- Why are the assumptions reasonable?
  - Normal distribution: the data are distributed approximately bell-shaped



# Remark - Model assumptions

*Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  ( $n = 16$ )  
(The data are realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )

**runtimes**




- Why are the assumptions reasonable?
  - Normal distribution: the data are distributed approximately bell-shaped
  - Knowledge of the standard deviation  $\sigma$ : Actually, this assumption is nonsense – why should we know something about the variance, when we do not know the expectation? Here assumed to be known for simplicity. Next lecture: replace  $\sigma$  by estimate  $s \rightarrow$  yields  $t$ -test

# Remark - Model assumptions

*Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  ( $n = 16$ )  
(The data are realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )

**runtimes**




- Why are the assumptions reasonable?
  - Normal distribution: the data are distributed approximately bell-shaped
  - Knowledge of the standard deviation  $\sigma$ : Actually, this assumption is nonsense – why should we know something about the variance, when we do not know the expectation? Here assumed to be known for simplicity. Next lecture: replace  $\sigma$  by estimate  $s \rightarrow$  yields  $t$ -test
  - Independence: We do not have a reason to assume that the observed individuals have too much in common.

# Remark - Model assumptions

*Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, 11^2)$  and  $\mu \in \mathbb{R}$  ( $n = 16$ )  
(The data are realizations of i.i.d. normal-distributed RVs, with unknown expectation  $\mu \in \mathbb{R}$ , but known variance  $\sigma^2 = 121$ )

**runtimes**



- Why are the assumptions reasonable?
  - Normal distribution: the data are distributed approximately bell-shaped
  - Knowledge of the standard deviation  $\sigma$ : Actually, this assumption is nonsense – why should we know something about the variance, when we do not know the expectation? Here assumed to be known for simplicity. Next lecture: replace  $\sigma$  by estimate  $s \rightarrow$  yields  $t$ -test
  - Independence: We do not have a reason to assume that the observed individuals have too much in common.
  - In general: A model is always a simplification. The description of the 'reality' through a theoretical construct ('model') is basically always 'wrong'. A complicated model (which is possibly not appropriately understood) is often useless. Models should be chosen as 'simple' objects.
  - George Box: 'All models are wrong' (but some are useful)

## Remark - Test statistic

The *test statistic* should accomplish two things:

1. It should measure the discrepancy between the **data** and the null hypothesis  $H_0$

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

Otherwise the procedure was nonsense.

## Remark - Test statistic

The *test statistic* should accomplish two things:

1. It should measure the discrepancy between the **data** and the null hypothesis  $H_0$

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$


Otherwise the procedure was nonsense.

2. It should be chosen such that its distribution was known under  $H_0$

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$

Otherwise we can not judge the discrepancy in 1.

Distribution of Z under  $H_0 : \mu = 30$



# Remark - Test statistic

The *test statistic* should accomplish two things:

1. It should measure the discrepancy between the **data** and the null hypothesis  $H_0$

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$


Otherwise the procedure was nonsense.

2. It should be chosen such that its distribution was known under  $H_0$

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$

Otherwise we can not judge the discrepancy in 1.

Distribution of Z under  $H_0 : \mu = 30$



- The abbreviation  $' \stackrel{H_0}{\sim}$  in  $Z \stackrel{H_0}{\sim} N(0, 1)$  means that the left hand side is distributed according to the right hand side, if the null hypothesis holds true, short: 'under  $H_0$ '

# Remark - Test statistic

The *test statistic* should accomplish two things:

1. It should measure the discrepancy between the **data** and the null hypothesis  $H_0$

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$


Otherwise the procedure was nonsense.

2. It should be chosen such that its distribution was known under  $H_0$

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$

Otherwise we can not judge the discrepancy in 1.

Distribution of Z under  $H_0 : \mu = 30$



- The abbreviation  $\stackrel{H_0}{\sim}$  in  $Z \stackrel{H_0}{\sim} N(0, 1)$  means that the left hand side is distributed according to the right hand side, if the null hypothesis holds true, short: 'under  $H_0$ '
- $z$  measures the discrepancy of  $\bar{x}$  from  $\mu_0$  in the units  $\sigma / \sqrt{n}$ , i.e., according to the variability of  $\bar{X}$ . Under  $H_0$ , the 'typical' deviation is one unit. In the data we observed  $|z| \approx 4.9$  units. This is untypically large!

# Remark - Test statistic

The *test statistic* should accomplish two things:

1. It should measure the discrepancy between the **data** and the null hypothesis  $H_0$

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}}$$

Otherwise the procedure was nonsense.

2. It should be chosen such that its distribution was known under  $H_0$

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}} \stackrel{H_0}{\sim} N(0, 1)$$

Otherwise we can not judge the discrepancy in 1.

- The abbreviation ' $\stackrel{H_0}{\sim}$ ' in  $Z \stackrel{H_0}{\sim} N(0, 1)$  means that the left hand side is distributed according to the right hand side, if the null hypothesis holds true, short: 'under  $H_0$ '
- $z$  measures the discrepancy of  $\bar{x}$  from  $\mu_0$  in the units  $\sigma / \sqrt{n}$ , i.e., according to the variability of  $\bar{X}$ . Under  $H_0$ , the 'typical' deviation is one unit. In the data we observed  $|z| \approx 4.9$  units. This is untypically large!
- If on the other hand  $H_0$  does not hold true, i.e., if the  $X_i$  have an expectation  $\mu_1$ , with  $\mu_1 \neq \mu_0$ , then  $Z$  is not distributed according to  $N(0, 1)$ , as we did not center correctly




## Remark - $p$ -value

$p$ -value (quantifies discrepancy)

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if the null hypothesis holds true.

Distribution of  $Z$  under  $H_0 : \mu = 30$




# Remark - $p$ -value

$p$ -value (quantifies discrepancy)

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if the null hypothesis holds true.

Distribution of  $Z$  under  $H_0 : \mu = 30$



- Index  $H_0$  again means : 'if the null hypothesis holds true'


# Remark - $p$ -value

$p$ -value (quantifies discrepancy)

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if the null hypothesis holds true.

Distribution of  $Z$  under  $H_0 : \mu = 30$



- Index  $H_0$  again means : 'if the null hypothesis holds true'
- $p$ -value 'small'  $\leftrightarrow$  If  $H_0$  holds true, then something unlikely has occurred  $\leftrightarrow$  incompatibility of the the **data** and the **assertion**.


# Remark - $p$ -value

$p$ -value (quantifies discrepancy)

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if the null hypothesis holds true.

Distribution of  $Z$  under  $H_0 : \mu = 30$



- Index  $H_0$  again means : 'if the null hypothesis holds true'
- $p$ -value 'small'  $\leftrightarrow$  If  $H_0$  holds true, then something unlikely has occurred  $\leftrightarrow$  incompatibility of the **data** and the **assertion**.
- The choice of 'small' / 'unlikely' is to be fixed by the statistician in advance(!). This is done by the choice of the *significance level*  $\alpha \in (0, 1)$ . Often  $\alpha = 5\%$


# Remark - $p$ -value

$p$ -value (quantifies discrepancy)

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if the null hypothesis holds true.

Distribution of  $Z$  under  $H_0 : \mu = 30$



- Index  $H_0$  again means : 'if the null hypothesis holds true'
- $p$ -value 'small'  $\leftrightarrow$  If  $H_0$  holds true, then something unlikely has occurred  $\leftrightarrow$  incompatibility of the the **data** and the **assertion**.
- The choice of 'small' / 'unlikely' is to be fixed by the statistician in advance(!). This is done by the choice of the *significance level*  $\alpha \in (0, 1)$ . Often  $\alpha = 5\%$
- Decision rule:
  - $p \leq \alpha \leftrightarrow$  incompatible enough  $\leftrightarrow$  reject null hypothesis


# Remark - $p$ -value

$p$ -value (quantifies discrepancy)

$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Probability to make an observation which is *at least as extreme* as in the data, if the null hypothesis holds true.

Distribution of  $Z$  under  $H_0 : \mu = 30$



- Index  $H_0$  again means : 'if the null hypothesis holds true'
- $p$ -value 'small'  $\leftrightarrow$  If  $H_0$  holds true, then something unlikely has occurred  $\leftrightarrow$  incompatibility of the the **data** and the **assertion**.
- The choice of 'small' / 'unlikely' is to be fixed by the statistician in advance(!). This is done by the choice of the *significance level*  $\alpha \in (0, 1)$ . Often  $\alpha = 5\%$
- Decision rule:
  - $p \leq \alpha \leftrightarrow$  incompatible enough  $\leftrightarrow$  reject null hypothesis
  - $p > \alpha \leftrightarrow$  not incompatible  $\leftrightarrow$  do *not* reject the null hypothesis

## Remark - $p$ -value and rejection area


- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- |   |  |
|---|--|
| <ul style="list-style-type: none"><li>• <math>p \leq \alpha</math></li><li>• <math>p &gt; \alpha</math></li></ul> | <ul style="list-style-type: none"><li><math>\leftrightarrow</math> reject <math>H_0</math></li><li><math>\leftrightarrow</math> do not reject <math>H_0</math></li></ul> |
|---|--|

## Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- |                   |                                       |
|-------------------|---------------------------------------|
| • $p \leq \alpha$ | $\leftrightarrow$ reject $H_0$        |
| • $p > \alpha$    | $\leftrightarrow$ do not reject $H_0$ |




- A different point of view of the upper decision rule: the *rejection area*  $R$



# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- |                   |                                       |
|-------------------|---------------------------------------|
| • $p \leq \alpha$ | $\leftrightarrow$ reject $H_0$        |
| • $p > \alpha$    | $\leftrightarrow$ do not reject $H_0$ |





- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- $p \leq \alpha \quad \leftrightarrow \text{reject } H_0$
- $p > \alpha \quad \leftrightarrow \text{do not reject } H_0$





- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- |                   |                                       |
|-------------------|---------------------------------------|
| • $p \leq \alpha$ | $\leftrightarrow$ reject $H_0$        |
| • $p > \alpha$    | $\leftrightarrow$ do not reject $H_0$ |




- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\leftrightarrow$  rejection at high discrepancy

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- |                   |                                       |
|-------------------|---------------------------------------|
| • $p \leq \alpha$ | $\leftrightarrow$ reject $H_0$        |
| • $p > \alpha$    | $\leftrightarrow$ do not reject $H_0$ |




- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\leftrightarrow$  rejection at high discrepancy
  - Set  $R$  such that  $\mathbb{P}_{H_0}(Z \in R) = \alpha$ . Here  $\alpha = 5\% \rightarrow R \approx (-\infty, -1.96] \cup [1.96, \infty)$   
Meaning: If  $H_0$  holds true, then we falsely reject with probability  $\alpha$

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- $p \leq \alpha \quad \leftrightarrow \text{reject } H_0$
- $p > \alpha \quad \leftrightarrow \text{do not reject } H_0$




- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\leftrightarrow$  rejection at high discrepancy
  - Set  $R$  such that  $\mathbb{P}_{H_0}(Z \in R) = \alpha$ . Here  $\alpha = 5\% \rightarrow R \approx (-\infty, -1.96] \cup [1.96, \infty)$   
Meaning: If  $H_0$  holds true, then we falsely reject with probability  $\alpha$
  - Equivalent to decision rule via  $p$ -value

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \leftrightarrow \text{reject } H_0$
- $p > \alpha \Leftrightarrow \text{do not reject } H_0$





- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\leftrightarrow$  rejection at high discrepancy
  - Set  $R$  such that  $\mathbb{P}_{H_0}(Z \in R) = \alpha$ . Here  $\alpha = 5\% \rightarrow R \approx (-\infty, -1.96] \cup [1.96, \infty)$   
Meaning: If  $H_0$  holds true, then we falsely reject with probability  $\alpha$
  - Equivalent to decision rule via  $p$ -value

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \leftrightarrow \text{reject } H_0$
- $p > \alpha \Leftrightarrow z \notin R \leftrightarrow \text{do not reject } H_0$





- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\leftrightarrow$  rejection at high discrepancy
  - Set  $R$  such that  $\mathbb{P}_{H_0}(Z \in R) = \alpha$ . Here  $\alpha = 5\% \rightarrow R \approx (-\infty, -1.96] \cup [1.96, \infty)$   
Meaning: If  $H_0$  holds true, then we falsely reject with probability  $\alpha$
  - Equivalent to decision rule via  $p$ -value

# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0$





- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\Leftrightarrow$  rejection at high discrepancy
  - Set  $R$  such that  $\mathbb{P}_{H_0}(Z \in R) = \alpha$ . Here  $\alpha = 5\% \rightarrow R \approx (-\infty, -1.96] \cup [1.96, \infty)$   
Meaning: If  $H_0$  holds true, then we falsely reject with probability  $\alpha$
  - Equivalent to decision rule via  $p$ -value



# Remark - $p$ -value and rejection area

- Significance level  $\alpha \in (0, 1)$
- $p$ -value:  $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$




- A different point of view of the upper decision rule: the *rejection area*  $R$ 
  - Reject  $H_0$  if and only if  $z \in R$
  - set  $R$  to the tails of the distribution  $\Leftrightarrow$  rejection at high discrepancy
  - Set  $R$  such that  $\mathbb{P}_{H_0}(Z \in R) = \alpha$ . Here  $\alpha = 5\% \rightarrow R \approx (-\infty, -1.96] \cup [1.96, \infty)$   
Meaning: If  $H_0$  holds true, then we falsely reject with probability  $\alpha$
  - Equivalent to decision rule via  $p$ -value

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$




- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$




- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$




- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$




- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$




- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$





- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting
- popular choices:  $\alpha = 5\%, 1\%, 0.1\%$ .

# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:

- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$





- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting
- popular choices:  $\alpha = 5\%, 1\%, 0.1\%$ .
- Here: Reject  $H_0$  on the 5% level, but not on the 0.1% level



# Remark - significance level

- Significance level  $\alpha \in (0, 1)$
- $p$ -Wert:  $P = \mathbb{P}_{H_0}(|Z| \geq |z|)$
- Decision rule:


- $p \leq \alpha \Leftrightarrow z \in R \Leftrightarrow \text{reject } H_0 \Leftrightarrow \text{say: 'the discrepancy was significant'}$
- $p > \alpha \Leftrightarrow z \notin R \Leftrightarrow \text{do not reject } H_0 \Leftrightarrow \text{'discrepancy was not significant'}$



- Decrease of  $\alpha \Leftrightarrow$  smaller rejection area  $R$ , i.e., more 'strict' with rejecting
- popular choices:  $\alpha = 5\%, 1\%, 0.1\%$ .
- Here: Reject  $H_0$  on the 5% level, but not on the 0.1% level
- Important: The level  $\alpha$  has to be chosen in advance! It is self-delusive to increase  $\alpha$  in order to reject!

## Remark - Interpretation

A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!" And further he claims: "The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."




$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

What is our opinion on that? (Apart from the fact, that we might find the delicate statement inappropriate from an interpersonal level)

## Remark - Interpretation

A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!" And further he claims: "The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

What is our opinion on that? (Apart from the fact, that we might find the delicate statement inappropriate from an interpersonal level)

# Remark - Interpretation

A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!" And further he claims: "The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

What is our opinion on that? (Apart from the fact, that we might find the delicate statement inappropriate from an interpersonal level)

- The data are not at all compatible with the assertion

# Remark - Interpretation

A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!" And further he claims: "The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

What is our opinion on that? (Apart from the fact, that we might find the delicate statement inappropriate from an interpersonal level)

- The **data** are **not at all compatible** with the **assertion**
- If the colleague is **right**, then a discrepancy which is at least as extreme as in the observed **data**, will occur in less than one of 1 million cases (as  $p \approx 9 \cdot 10^{-7} < 10^{-6}$ )

# Remark - Interpretation

A skeptic colleague claims: "the course is useless. The 16 students were just over average beforehand!" And further he claims: "The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been  $\mu_0 = 30$ ."





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

What is our opinion on that? (Apart from the fact, that we might find the delicate statement inappropriate from an interpersonal level)

- The **data** are **not at all compatible** with the **assertion**
- If the colleague is **right**, then a discrepancy which is at least as extreme as in the observed **data**, will occur in less than one of 1 million cases (as  $p \approx 9 \cdot 10^{-7} < 10^{-6}$ )
- → The **data** give us good reason to **doubt** the **assertion**!

# Remark - Erroneous interpretation





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:

- the colleague is right, respectively the null hypothesis holds true

# Remark - Erroneous interpretation





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:

- the colleague is right, respectively the null hypothesis holds true



# Remark - Erroneous interpretation





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:

- ~~the colleague is right, respectively the null hypothesis holds true~~
- ~~the colleague is wrong, resp. the null hypothesis does not hold true~~

# Remark - Erroneous interpretation





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:

- ~~the colleague is right, respectively the null hypothesis holds true~~
- ~~the colleague is wrong, resp. the null hypothesis does not hold true~~
- ~~the colleague is probably right / wrong, resp. the null hypothesis is probably true / false~~

# Remark - Erroneous interpretation





$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:

- ~~the colleague is right, respectively the null hypothesis holds true~~
- ~~the colleague is wrong, resp. the null hypothesis does not hold true~~
- ~~the colleague is probably right / wrong, resp. the null hypothesis is probably true / false~~

We cannot say that, because we just make **probability statements** about the **data**, in fact under the theoretical assumption that the **null hypothesis** holds true.

# Remark - Erroneous interpretation



$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$



Popular misinterpretations:

- ~~the colleague is right, respectively the null hypothesis holds true~~
- ~~the colleague is wrong, resp. the null hypothesis does not hold true~~
- ~~the colleague is probably right / wrong, resp. the null hypothesis is probably true / false~~

We cannot say that, because we just make **probability statements** about the **data**, in fact under the theoretical assumption that the **null hypothesis** holds true.

We cannot say anything about the huge unknown population / the null hypothesis!

# Remark - Erroneous interpretation



$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:



- ~~the colleague is right, respectively the null hypothesis holds true~~
- ~~the colleague is wrong, resp. the null hypothesis does not hold true~~
- ~~the colleague is probably right / wrong, resp. the null hypothesis is probably true / false~~

We cannot say that, because we just make **probability statements** about the **data**, in fact under the theoretical assumption that the **null hypothesis** holds true.

We cannot say anything about the huge unknown population / the null hypothesis!

Rejecting the null hypothesis does not mean that its wrong or probably wrong!

# Remark - Erroneous interpretation



$$p = \mathbb{P}_{H_0}(|Z| \geq |z|) \approx 9 \cdot 10^{-7}$$

Popular misinterpretations:

- ~~the colleague is right, respectively the null hypothesis holds true~~
- ~~the colleague is wrong, resp. the null hypothesis does not hold true~~
- ~~the colleague is probably right / wrong, resp. the null hypothesis is probably true / false~~



We cannot say that, because we just make **probability statements** about the **data**, in fact under the theoretical assumption that the **null hypothesis** holds true.

We cannot say anything about the huge unknown population / the null hypothesis!

Rejecting the null hypothesis does not mean that its wrong or probably wrong!



Throughout the course we will stay in the realm of the so-called *frequentists approach* of statistics, where the unknown parameters (like  $\mu$ ) are treated as *non-random*. In contrast, the *Bayesian* statisticians will want to make probability statements about the parameter as in their world parameters are modeled as random variables.

# Two-sided and one-sided testing



- $H_0$  rejected  $\leftrightarrow z \in R$



# Two-sided and one-sided testing



- $H_0$  rejected  $\leftrightarrow z \in R$
- Let  $q_\alpha$  be the  $\alpha$ -quantile of  $N(0, 1)$ , i.e.,  $\mathbb{P}(Y < q_\alpha) = \alpha$  for  $Y \sim N(0, 1)$





# Two-sided and one-sided testing





- $H_0$  rejected  $\leftrightarrow z \in R$
- Let  $q_\alpha$  be the  $\alpha$ -quantile of  $N(0, 1)$ , i.e.,  $\mathbb{P}(Y < q_\alpha) = \alpha$  for  $Y \sim N(0, 1)$
- Differentiation between *two-sided* and *one-sided* tests

# Two-sided and one-sided testing





- $H_0$  rejected  $\leftrightarrow z \in R$
- Let  $q_\alpha$  be the  $\alpha$ -quantile of  $N(0, 1)$ , i.e.,  $\mathbb{P}(Y < q_\alpha) = \alpha$  for  $Y \sim N(0, 1)$
- Differentiation between *two-sided* and *one-sided* tests
  - two-sided: null hypothesis  $H_0 : \mu = \mu_0$  ( $\rightarrow$  alternative  $H_A : \mu \neq \mu_0$ )  
 $R = (-\infty, q_{\alpha/2}] \cup [q_{1-\alpha/2}, \infty)$   
 $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$   
Extreme values if  $z$  speak against the null hypothesis

# Two-sided and one-sided testing





- $H_0$  rejected  $\leftrightarrow z \in R$
- Let  $q_\alpha$  be the  $\alpha$ -quantile of  $N(0, 1)$ , i.e.,  $\mathbb{P}(Y < q_\alpha) = \alpha$  for  $Y \sim N(0, 1)$
- Differentiation between *two-sided* and *one-sided* tests
  - two-sided: null hypothesis  $H_0 : \mu = \mu_0$  ( $\rightarrow$  alternative  $H_A : \mu \neq \mu_0$ )  
 $R = (-\infty, q_{\alpha/2}] \cup [q_{1-\alpha/2}, \infty)$   
 $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$   
Extreme values if  $z$  speak against the null hypothesis
  - left-sided: null hypothesis  $H_0 : \mu \geq \mu_0$  ( $\rightarrow$  alternative  $H_A : \mu < \mu_0$ )  
 $R = (-\infty, q_\alpha]$   
 $p = \mathbb{P}_{H_0}(Z \leq z)$   
Small values of  $z$  speak against the null hypothesis

# Two-sided and one-sided testing





- $H_0$  rejected  $\leftrightarrow z \in R$
- Let  $q_\alpha$  be the  $\alpha$ -quantile of  $N(0, 1)$ , i.e.,  $\mathbb{P}(Y < q_\alpha) = \alpha$  for  $Y \sim N(0, 1)$
- Differentiation between *two-sided* and *one-sided* tests
  - two-sided: null hypothesis  $H_0 : \mu = \mu_0$  ( $\rightarrow$  alternative  $H_A : \mu \neq \mu_0$ )  
 $R = (-\infty, q_{\alpha/2}] \cup [q_{1-\alpha/2}, \infty)$   
 $p = \mathbb{P}_{H_0}(|Z| \geq |z|)$   
Extreme values if  $z$  speak against the null hypothesis
  - left-sided: null hypothesis  $H_0 : \mu \geq \mu_0$  ( $\rightarrow$  alternative  $H_A : \mu < \mu_0$ )  
 $R = (-\infty, q_\alpha]$   
 $p = \mathbb{P}_{H_0}(Z \leq z)$   
Small values of  $z$  speak against the null hypothesis
  - right sided: null hypothesis  $H_0 : \mu \leq \mu_0$  ( $\rightarrow$  alternative  $H_A : \mu > \mu_0$ )  
 $R = [q_{1-\alpha}, \infty)$   
 $p = \mathbb{P}_{H_0}(Z \geq z)$   
Large values of  $z$  speak against the null hypothesis

# Two-sided and one-sided testing





- In our example we could also have tested left-sided, because we knew in advance that the mean of the observed runtimes was smaller than the assertion, and we wanted to detect this fact of being 'faster'.

# Two-sided and one-sided testing





- In our example we could also have tested left-sided, because we knew in advance that the mean of the observed runtimes was smaller than the assertion, and we wanted to detect this fact of being 'faster'.
- However, we preferably formulate hypothesis before we observe the data.

# Two-sided and one-sided testing



- In our example we could also have tested left-sided, because we knew in advance that the mean of the observed runtimes was smaller than the assertion, and we wanted to detect this fact of being 'faster'.
- However, we preferably formulate hypothesis before we observe the data.
- And it might have happened that the mean observed runtime was slower (larger) than the assertion. This would then have suggested a 'negative effect', which would not have been detected with a left-sided test.

# Two-sided and one-sided testing




- In our example we could also have tested left-sided, because we knew in advance that the mean of the observed runtimes was smaller than the assertion, and we wanted to detect this fact of being 'faster'.
- However, we preferably formulate hypothesis before we observe the data.
- And it might have happened that the mean observed runtime was slower (larger) than the assertion. This would then have suggested a 'negative effect', which would not have been detected with a left-sided test.
- Thus, depending on the context we need to decide of whether to perform a one- or two-sided test. (Rule of thumb: two-sided)





# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true		



# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	





$H_0$  falsely rejected,  $\alpha$ -error committed

- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true

# Errors and test power



Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	



- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$

# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )




$H_0$  rightly NOT rejected

- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$

# Errors and test power



Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )
does not hold true		



- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$

# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )
does not hold true		$\beta$ -error ( $= \beta$ )





$H_0$  falsely NOT rejected,  $\beta$ -error committed

- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$
- $\beta$ -error:  $H_0$  is not rejected although  $H_0$  does not hold true

# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )
does not hold true		$\beta$ -error ( $= \beta$ )





$H_0$  falsely NOT rejected,  $\beta$ -error committed

- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$
- $\beta$ -error:  $H_0$  is not rejected although  $H_0$  does not hold true
- The probability  $\beta$  to commit the  $\beta$ -error depends on the concrete alternative

# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )
does not hold true		$\beta$ -error ( $= \beta$ )





- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in \mathbf{R}) = \alpha$
- $\beta$ -error:  $H_0$  is not rejected although  $H_0$  does not hold true
- The probability  $\beta$  to commit the  $\beta$ -error depends on the concrete alternative



# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )
does not hold true	(test power $= 1 - \beta$ )	$\beta$ -error ( $= \beta$ )





$H_0$  rightly rejected, test power  $1 - \beta$

- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$
- $\beta$ -error:  $H_0$  is not rejected although  $H_0$  does not hold true
- The probability  $\beta$  to commit the  $\beta$ -error depends on the concrete alternative
- Test power  $1 - \beta$  is the probability to reject  $H_0$ , if  $H_0$  does not hold true

# Errors and test power

Null hypothesis	rejected (with prob)	not rejected (with prob)
holds true	$\alpha$ -error ( $= \alpha$ )	( $= 1 - \alpha$ )
does not hold true	(test power $= 1 - \beta$ )	$\beta$ -error ( $= \beta$ )



$H_0$  rightly rejected, test power  $1 - \beta$

- $\alpha$ -error:  $H_0$  is rejected although  $H_0$  holds true
- The probability to commit the  $\alpha$ -error is given through the choice of the significance level  $\alpha$ , as by construction  $\mathbb{P}_{H_0}(Z \in R) = \alpha$
- $\beta$ -error:  $H_0$  is not rejected although  $H_0$  does not hold true
- The probability  $\beta$  to commit the  $\beta$ -error depends on the concrete alternative
- Test power  $1 - \beta$  is the probability to reject  $H_0$ , if  $H_0$  does not hold true

The question of whether we commit these errors can never be answered in practice, because hypotheses are theoretical assumptions

Thank you!