


Analysis of variance and multiple testing



All examples are fictitious. All data are simulated and the graphics were created with the statistical program package R.


The materials are protected by copyright and are only provided for personal use for studies at TU Vienna. Further use is not permitted. In particular, it is not permitted to distribute the materials or make them publicly available (e.g. in social networks, on learning platforms, etc.).

Sämtliche Beispiele sind frei erfunden. Alle Daten sind simuliert und die Grafiken wurden mit statistischen Programmpaket R erstellt.


Die Materialien sind urheberrechtlich geschützt und dürfen ausschließlich für den Eigengebrauch im Rahmen des Studiums an der TU Wien genutzt werden. Eine weitere Nutzung ist nicht gestattet. Insbesondere ist es nicht gestattet, die Materialien zu verbreiten oder öffentlich zugänglich zu machen (etwa im Rahmen sozialer Netzwerke, Lernplattformen etc.).

Reminder

At four universities students of a certain study program were interviewed regarding the level of their satisfaction with the study situation. An extensive survey had to be filled out. Subsequently, for every respondent a global value of 'contentment' was evaluated.




Reminder: two-sample situation




- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$

Reminder: two-sample situation




- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?

Reminder: two-sample situation




- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$

Reminder: two-sample situation





- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
- In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
$$X_i \sim N(\mu_1, \sigma_1^2) \text{ for } i = 1, \dots, n_1 \quad \text{and} \quad Y_j \sim N(\mu_2, \sigma_2^2) \text{ for } j = 1, \dots, n_2$$
with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$

Reminder: two-sample situation





- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
- In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
$$X_i \sim N(\mu_1, \sigma_1^2) \text{ for } i = 1, \dots, n_1 \quad \text{and} \quad Y_j \sim N(\mu_2, \sigma_2^2) \text{ for } j = 1, \dots, n_2$$
with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$ it holds under $H_0 : \mu_y - \mu_x = d_0$ that

Reminder: two-sample situation





- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
 - Mean \bar{x} and \bar{y} far apart?
 - Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
 - In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
 - $X_i \sim N(\mu_1, \sigma_1^2)$ for $i = 1, \dots, n_1$ and
 - $Y_j \sim N(\mu_2, \sigma_2^2)$ for $j = 1, \dots, n_2$ with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
- it holds under $H_0 : \mu_y - \mu_x = d_0$ that $T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$ (R knows ν)

Reminder: two-sample situation





- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
- In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
 $X_i \sim N(\mu_1, \sigma_1^2)$ for $i = 1, \dots, n_1$ and
 $Y_j \sim N(\mu_2, \sigma_2^2)$ for $j = 1, \dots, n_2$ with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
 it holds under $H_0 : \mu_y - \mu_x = d_0$ that $T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$ (R knows ν)
- Judge data via $t = \frac{(\bar{y} - \bar{x}) - d_0}{\sqrt{sem_y^2 + sem_x^2}}$ according to the $t(\nu)$ -distribution

Reminder: two-sample situation





- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
- In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
 $X_i \sim N(\mu_1, \sigma_1^2)$ for $i = 1, \dots, n_1$ and
 $Y_j \sim N(\mu_2, \sigma_2^2)$ for $j = 1, \dots, n_2$ with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
 it holds under $H_0 : \mu_y - \mu_x = d_0$ that $T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$ (R knows ν)
- Judge data via $t = \frac{(\bar{y} - \bar{x}) - d_0}{\sqrt{sem_y^2 + sem_x^2}}$ according to the $t(\nu)$ -distribution
- Example: for $d_0 = 5$ it is $p > \alpha = 5\%$ resp. $t \notin R$, thus do not reject H_0 on the α -level

Reminder: two-sample situation



- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
- In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
 $X_i \sim N(\mu_1, \sigma_1^2)$ for $i = 1, \dots, n_1$ and
 $Y_j \sim N(\mu_2, \sigma_2^2)$ for $j = 1, \dots, n_2$ with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
 it holds under $H_0 : \mu_y - \mu_x = d_0$ that $T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$ (R knows ν)
- Judge data via $t = \frac{(\bar{y} - \bar{x}) - d_0}{\sqrt{sem_y^2 + sem_x^2}}$ according to the $t(\nu)$ -distribution
- Example: for $d_0 = 5$ it is $p > \alpha = 5\%$ resp. $t \notin R$, thus do not reject H_0 on the α -level
- In practice often $d_0 = 0$, i.e., 'no shift' \rightarrow here rejected



Reminder: two-sample situation



- Data $x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}$
- Mean \bar{x} and \bar{y} far apart?
- Depends on the standard errors $sem_x = s_x / \sqrt{n_1}$ and $sem_y = s_y / \sqrt{n_2}$
- In the model $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$ independent RVs with
 $X_i \sim N(\mu_1, \sigma_1^2)$ for $i = 1, \dots, n_1$ and
 $Y_j \sim N(\mu_2, \sigma_2^2)$ for $j = 1, \dots, n_2$ with $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
 it holds under $H_0 : \mu_y - \mu_x = d_0$ that $T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$ (R knows ν)
- Judge data via $t = \frac{(\bar{y} - \bar{x}) - d_0}{\sqrt{sem_y^2 + sem_x^2}}$ according to the $t(\nu)$ -distribution
- Example: for $d_0 = 5$ it is $p > \alpha = 5\%$ resp. $t \notin R$, thus do not reject H_0 on the α -level
- In practice often $d_0 = 0$, i.e., 'no shift' \rightarrow here rejected


This is the Welch-test. Remember that there is also Student's t -test, where equal variances $\sigma_1^2 = \sigma_2^2$ are assumed, for which we found under H_0 that $T \sim t(n_1 + n_2 - 2)$ while a different scaling was used in T ('pooled variance')

Message




The t -statistic says:
the **discrepancy between the groups** is large,
if the **variability within the groups** is small

Today: Comparison of $k \geq 2$ groups



Here $k = 4$ groups

Today: Comparison of $k \geq 2$ groups




Here $k = 4$ groups

Two possibilities to deal with that:

- Multiple pairwise t -tests

Today: Comparison of $k \geq 2$ groups




Here $k = 4$ groups

Two possibilities to deal with that:

- Multiple pairwise t -tests
- Analysis of variance (ANOVA)


Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test



Multiple testing


Multiple pairwise t -tests: for every pair a two-sample t -test



● means

Multiple testing


Multiple pairwise t -tests: for every pair a two-sample t -test



● means \pm standard errors

Multiple testing


Multiple pairwise t -tests: for every pair a two-sample t -test



- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large

Multiple testing


Multiple pairwise t -tests: for every pair a two-sample t -test



- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test




- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p-value	red	green	green	green	red	red

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test




- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value	0.004					

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test




- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p-value	0.004	0.067				

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test




- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value	0.004	0.067	0.710			

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test




- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value	0.004	0.067	0.710	0.090		

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test




- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value	0.004	0.067	0.710	0.090	0.001	

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

Multiple pairwise t -tests: for every pair a two-sample t -test



- means \pm standard errors
- Naive: Discrepancy between A and D small, between B and D large
- $k = 4$ groups \rightarrow six t -tests ($d_0 = 0$, no shift)

groups	A/B	A/C	A/D	B/C	B/D	C/D
p-value	0.004	0.067	0.710	0.090	0.001	0.014

red: $p < 5\%$, green: $p \geq 5\%$

Multiple testing

- With 4 groups the number of t -tests performed is 6

Multiple testing

- With 4 groups the number of t -tests performed is 6
- How many t -tests do we have to perform with k groups?

Multiple testing

- With 4 groups the number of t -tests performed is 6
- How many t -tests do we have to perform with k groups? Answer:

$$m = \binom{k}{2}$$

(the number of 2-element subsets of a k -element set)

Multiple testing

- With 4 groups the number of t -tests performed is 6
- How many t -tests do we have to perform with k groups? Answer:

$$m = \binom{k}{2}$$

(the number of 2-element subsets of a k -element set)

- We mention that

$$m = \frac{k(k-1)}{2}$$

grows quadratically in k

Multiple testing

- With 4 groups the number of t -tests performed is 6
- How many t -tests do we have to perform with k groups? Answer:

$$m = \binom{k}{2}$$

(the number of 2-element subsets of a k -element set)

- We mention that

$$m = \frac{k(k-1)}{2}$$

grows quadratically in k

- Meaning: the number of tests to perform increases rapidly with the number of groups

Multiple testing

- With 4 groups the number of t -tests performed is 6
- How many t -tests do we have to perform with k groups? Answer:

$$m = \binom{k}{2}$$

(the number of 2-element subsets of a k -element set)

- We mention that

$$m = \frac{k(k-1)}{2}$$

grows quadratically in k

- Meaning: the number of tests to perform increases rapidly with the number of groups
- ... and the application of many statistical tests brings a conceptual problem ...

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true

The multiple testing problem


gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests?**

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$
- Message: The **probability to commit the α -error at least once** tended to 1, for $m \rightarrow \infty$ (' α -error cumulation')

The multiple testing problem

gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$
- Message: The **probability to commit the α -error at least once** tended to 1, for $m \rightarrow \infty$ (' α -error cumulation')
- And this is bad news! If we only perform enough tests, then the null hypothesis will be rejected at some point (even if it was always true)

The multiple testing problem


gedankenexperiment

- assume we performed m (independent) statistical tests at level α
- and we knew, that the null hypothesis H_0 was true
- What was the **probability** to commit the α -error (i.e., falsely reject) **in at least one of the tests**?




- In the picture $\alpha = 5\%$
- Message: The **probability to commit the α -error at least once** tended to 1, for $m \rightarrow \infty$ (' α -error cumulation')
- And this is bad news! If we only perform enough tests, then the null hypothesis will be rejected at some point (even if it was always true)
- Meaning for practice: do not perform unnecessary tests!

The multiple testing problem




- m (independent) tests at level α
- Assume H_0 always holds true
- Looking for the **probability to commit the α -error at least once**

The multiple testing problem



- m (independent) tests at level α
- Assume H_0 always holds true
- Looking for the **probability to commit the α -error at least once**
- Let V_i be the event, that the i -th null hypothesis is rejected

The multiple testing problem




- m (independent) tests at level α
- Assume H_0 always holds true
- Looking for the **probability to commit the α -error at least once**
- Let V_i be the event, that the i -th null hypothesis is rejected
 - $m = 1$

$$\mathbb{P}_{H_0}(V_1) = \alpha$$

by construction of the test

The multiple testing problem



- m (independent) tests at level α
- Assume H_0 always holds true
- Looking for the **probability to commit the α -error at least once**
- Let V_i be the event, that the i -th null hypothesis is rejected
 - $m = 1$

$$\mathbb{P}_{H_0}(V_1) = \alpha$$


by construction of the test

- $m = 2$:

$$\mathbb{P}_{H_0}(V_1 \cup V_2) = 1 - \mathbb{P}_{H_0}(V_1^c \cap V_2^c) = 1 - \mathbb{P}_{H_0}(V_1^c)\mathbb{P}_{H_0}(V_2^c) = 1 - (1 - \alpha)^2$$

used: $\mathbb{P}(\text{'at least one'}) = 1 - \mathbb{P}(\text{'none'})$ as well as independence

The multiple testing problem



- m (independent) tests at level α
- Assume H_0 always holds true
- Looking for the **probability to commit the α -error at least once**
- Let V_i be the event, that the i -th null hypothesis is rejected
 - $m = 1$

$$\mathbb{P}_{H_0}(V_1) = \alpha$$

by construction of the test

- $m = 2$:


$$\mathbb{P}_{H_0}(V_1 \cup V_2) = 1 - \mathbb{P}_{H_0}(V_1^c \cap V_2^c) = 1 - \mathbb{P}_{H_0}(V_1^c)\mathbb{P}_{H_0}(V_2^c) = 1 - (1 - \alpha)^2$$

used: $\mathbb{P}(\text{'at least one'}) = 1 - \mathbb{P}(\text{'none'})$ as well as independence

- general m :

$$\mathbb{P}_{H_0}\left(\bigcup_{i=1}^m V_i\right) = 1 - \mathbb{P}_{H_0}\left(\bigcap_{i=1}^m V_i^c\right) = 1 - \prod_{i=1}^m \mathbb{P}_{H_0}(V_i^c) = \boxed{1 - (1 - \alpha)^m}$$

The multiple testing problem



- m (independent) tests at level α
- Assume H_0 always holds true
- Looking for the **probability to commit the α -error at least once**
- Let V_i be the event, that the i -th null hypothesis is rejected
 - $m = 1$

$$\mathbb{P}_{H_0}(V_1) = \alpha$$

by construction of the test

- $m = 2$:

$$\mathbb{P}_{H_0}(V_1 \cup V_2) = 1 - \mathbb{P}_{H_0}(V_1^c \cap V_2^c) = 1 - \mathbb{P}_{H_0}(V_1^c)\mathbb{P}_{H_0}(V_2^c) = 1 - (1 - \alpha)^2$$

used: $\mathbb{P}(\text{'at least one'}) = 1 - \mathbb{P}(\text{'none'})$ as well as independence

- general m :

$$\mathbb{P}_{H_0}\left(\bigcup_{i=1}^m V_i\right) = 1 - \mathbb{P}_{H_0}\left(\bigcap_{i=1}^m V_i^c\right) = 1 - \prod_{i=1}^m \mathbb{P}_{H_0}(V_i^c) = \boxed{1 - (1 - \alpha)^m} \rightarrow 1 \quad (m \rightarrow \infty)$$

Way out: Bonferroni correction

- m tests at level α
- Problem: the global α -error is increased

Way out: Bonferroni correction

- m tests at level α
- Problem: the global α -error is increased
- A possible way out: be more 'strict' with each of the m tests

Way out: Bonferroni correction

- m tests at level α
- Problem: the **global** α -error is increased
- A possible way out: be more 'strict' with each of the m tests
- More precisely: Correct the original significance level α to

$$\alpha^* = \frac{\alpha}{m}$$

known as the *Bonferroni-correction*

('divide by number of tests')

Way out: Bonferroni correction

- m tests at level α
- Problem: the **global** α -error is increased
- A possible way out: be more 'strict' with each of the m tests
- More precisely: Correct the original significance level α to

$$\alpha^* = \frac{\alpha}{m}$$

known as the *Bonferroni-correction*

('divide by number of tests')

- Consequence: the probability **to commit the α -alpha error at least once** is restricted by α

Way out: Bonferroni correction

- m tests at level α
- Problem: the **global** α -error is increased
- A possible way out: be more 'strict' with each of the m tests
- More precisely: Correct the original significance level α to

$$\alpha^* = \frac{\alpha}{m}$$

known as the *Bonferroni-correction*

('divide by number of tests')

- Consequence: the probability **to commit the α -alpha error at least once** is restricted by α
- Quantitatively:

$$\mathbb{P}_{H_0}\left(\bigcup_{i=1}^m V_i\right) \leq \sum_{i=1}^m \mathbb{P}_{H_0}(V_i) = m \cdot \alpha^* = \alpha$$

Way out: Bonferroni correction

- m tests at level α
- Problem: the **global** α -error is increased
- A possible way out: be more 'strict' with each of the m tests
- More precisely: Correct the original significance level α to

$$\alpha^* = \frac{\alpha}{m}$$

known as the *Bonferroni-correction*

(‘divide by number of tests’)

- Consequence: the probability **to commit the α -alpha error at least once** is restricted by α
- Quantitatively:

$$\mathbb{P}_{H_0}\left(\bigcup_{i=1}^m V_i\right) \leq \sum_{i=1}^m \mathbb{P}_{H_0}(V_i) = m \cdot \alpha^* = \alpha$$

- Alternatively: Use the original level α , but correct every p -value p to

$$p^* = m \cdot p$$

(‘multiplication with the number of tests’)

Way out: Bonferroni correction

- m tests at level α
- Problem: the **global** α -error is increased
- A possible way out: be more 'strict' with each of the m tests
- More precisely: Correct the original significance level α to

$$\alpha^* = \frac{\alpha}{m}$$

known as the *Bonferroni-correction*

('divide by number of tests')

- Consequence: the probability **to commit the α -alpha error at least once** is restricted by α
- Quantitatively:

$$\mathbb{P}_{H_0}\left(\bigcup_{i=1}^m V_i\right) \leq \sum_{i=1}^m \mathbb{P}_{H_0}(V_i) = m \cdot \alpha^* = \alpha$$


- Alternatively: Use the original level α , but correct every p -value p to

$$p^* = m \cdot p$$

('multiplication with the number of tests')

- Equivalent, as $p \leq \alpha^* \Leftrightarrow p^* \leq \alpha$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)						

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024					

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268				

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268	1			

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268	1	0.540		

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268	1	0.540	0.006	

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268	1	0.540	0.006	0.084

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268	1	0.540	0.006	0.084

red: $< 5\%$, green: $\geq 5\%$

- Bonferroni: multiply the original p -value with $m = 6$
- One additional null hypothesis cannot be rejected after the Bonferroni correction.

Multiple testing and Bonferroni-correction




groups	A/B	A/C	A/D	B/C	B/D	C/D
p -value p (prior Bonferroni)	0.004	0.067	0.710	0.090	0.001	0.014
p -value p^* (after Bonferroni)	0.024	0.268	1	0.540	0.006	0.084

red: $< 5\%$, green: $\geq 5\%$


- Bonferroni: multiply the original p -value with $m = 6$
- One additional null hypothesis cannot be rejected after the Bonferroni correction. We are more strict!

Analysis of variance



- $k = 4$ groups
- $m = 6$ pairwise t -tests \rightarrow the global α -error is increased

Analysis of variance




- $k = 4$ groups
- $m = 6$ pairwise t -tests \rightarrow the global α -error is increased
- Another possible 'way out':

The analysis of variance

short ANOVA

Analysis of variance




- $k = 4$ groups
- $m = 6$ pairwise t -tests \rightarrow the global α -error is increased
- Another possible 'way out':

The analysis of variance

short ANOVA

- Basic idea: test all k groups simultaneously

Analysis of variance




- $k = 4$ groups
- $m = 6$ pairwise t -tests \rightarrow the global α -error is increased
- Another possible 'way out':

The analysis of variance

short ANOVA

- Basic idea: test all k groups simultaneously
- Null hypothesis: the data all derive from the same normal distribution (\rightarrow particularly no shift)

Analysis of variance



- $k = 4$ groups
- $m = 6$ pairwise t -tests \rightarrow the global α -error is increased
- Another possible 'way out':

The analysis of variance

short ANOVA

- Basic idea: test all k groups simultaneously
- Null hypothesis: the data all derive from the same normal distribution (\rightarrow particularly no shift)
- Intuitively in the graphic: this is not very plausible, because the **group means** are partly **far away (multiple sem)** from each other

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



• Notation

- k groups

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



• Notation

- k groups
- n_i observations in the i -th group

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



Notation

- k groups
- n_i observations in the i -th group
- the j -th observation in the i -th group is $X_{i,j}$

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



- Notation
 - k groups
 - n_i observations in the i -th group
 - the j -th observation in the i -th group is $X_{i,j}$
- Assumptions
 - independence

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



- Notation
 - k groups
 - n_i observations in the i -th group
 - the j -th observation in the i -th group is $X_{i,j}$
- Assumptions
 - independence
 - normal distribution

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



- Notation
 - k groups
 - n_i observations in the i -th group
 - the j -th observation in the i -th group is $X_{i,j}$
- Assumptions
 - independence
 - normal distribution
 - own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$




- Notation
 - k groups
 - n_i observations in the i -th group
 - the j -th observation in the i -th group is $X_{i,j}$
- Assumptions
 - independence
 - normal distribution
 - own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)
 - the variance σ^2 is constant

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



• Notation

- k groups
- n_i observations in the i -th group
- the j -th observation in the i -th group is $X_{i,j}$


• Assumptions

- independence
- normal distribution
- own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)
- the variance σ^2 is constant

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$




- Notation
 - k groups
 - n_i observations in the i -th group
 - the j -th observation in the i -th group is $X_{i,j}$
- Assumptions
 - independence
 - normal distribution
 - own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)
 - the variance σ^2 is constant

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



• Notation

- k groups
- n_i observations in the i -th group
- the j -th observation in the i -th group is $X_{i,j}$


• Assumptions

- independence
- normal distribution
- own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)
- the variance σ^2 is constant

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



• Notation

- k groups
- n_i observations in the i -th group
- the j -th observation in the i -th group is $X_{i,j}$


• Assumptions

- independence
- normal distribution
- own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)
- the variance σ^2 is constant

ANOVA - Model

Model:

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$




- Notation
 - k groups
 - n_i observations in the i -th group
 - the j -th observation in the i -th group is $X_{i,j}$
- Assumptions
 - independence
 - normal distribution
 - own (unknown) expectation μ_i in the i -th group (i.e., possible 'shift' between groups)
 - the variance σ^2 is constant

Note that for $k = 2$ groups, this model equals the model in Student's two-sample t -test.

ANOVA - Null hypothesis


Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



- Null hypothesis: $H_0 : \mu_1 = \mu_2 = \dots = \mu_k$
the distributions are equal, in particular no shift

ANOVA - Null hypothesis


Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



- Null hypothesis: $H_0 : \mu_1 = \mu_2 = \dots = \mu_k$
the distributions are equal, in particular no shift
- Alternative: at least two expectations are not equal, $\mu_i \neq \mu_\ell$



ANOVA - Null hypothesis

Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$





- Null hypothesis: $H_0 : \mu_1 = \mu_2 = \dots = \mu_k$
the distributions are equal, in particular no shift
- Alternative: at least two expectations are not equal, $\mu_i \neq \mu_\ell$
- How can we test the null hypothesis?

Reminder: t -Statistic



The t -statistic says:
the **discrepancy between the groups** is large,
if the **variability within the groups** is small

Idea of the ANOVA analog




The f -statistic will tell us:
the **variability between the groups** is large,
if the **variability within the groups** is small

F-Statistic

Naive:


$$f \stackrel{!}{=} \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$



F-Statistic

Naive:

$$f \stackrel{!}{=} \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$




- Need 1.: group means, for $i = 1, \dots, k$

$$\bar{X}_{i,\cdot} := \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$$

F-Statistic

Naive:

$$f \stackrel{!}{=} \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$



- Need 1.: group means, for $i = 1, \dots, k$


$$\bar{X}_{i,\cdot} := \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$$

- Need 2.: global mean over all groups

$$\bar{X} := \frac{1}{n} \sum_{i,j} X_{i,j}$$


with $n = n_1 + \dots + n_k$ total number of observations

F-Statistic (google: R. A. Fisher)



- $\bar{X}_{i.} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$


F-Statistic (google: R. A. Fisher)



- $\bar{X}_{i,\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F -Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2}$$


F-Statistic (google: R. A. Fisher)



- $\bar{X}_{i,\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F -Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

F-Statistic (google: R. A. Fisher)




- $\bar{X}_{i,\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F -Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

- $F \geq 0$, due to squaring

F-Statistic (google: R. A. Fisher)




- $\bar{X}_{i,\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F-Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

- $F \geq 0$, due to squaring
- If the evaluation of the data f is very large, this speaks against H_0

F-Statistic (google: R. A. Fisher)




- $\bar{X}_{i,\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F -Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

- $F \geq 0$, due to squaring
- If the evaluation of the data f is very large, this speaks against H_0
 - Because then the **numerator** is large compared to the **denominator**

F-Statistic (google: R. A. Fisher)




- $\bar{X}_{i,\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F -Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

- $F \geq 0$, due to squaring
- If the evaluation of the data f is very large, this speaks against H_0
 - Because then the **numerator** is large compared to the **denominator**
 - The test is always right-sided (only large values speak against H_0)

F-Statistic (google: R. A. Fisher)




- $\bar{X}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F-Statistic is

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

- $F \geq 0$, due to squaring
- If the evaluation of the data f is very large, this speaks against H_0
 - Because then the **numerator** is large compared to the **denominator**
 - The test is always right-sided (only large values speak against H_0)
- The distribution of F is known under H_0 : the Fisher-distribution with $(k-1)$ -numerator- and $(n-k)$ -denominator- degrees of freedom, short $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$

F-Statistic (google: R. A. Fisher)



- $\bar{X}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{i,j}$ and $\bar{X} = \frac{1}{n} \sum_{i,j} X_{i,j}$ with $n = \sum_{i=1}^k n_i$
- The F-Statistic is


$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$

- $F \geq 0$, due to squaring
- If the evaluation of the data f is very large, this speaks against H_0
 - Because then the **numerator** is large compared to the **denominator**
 - The test is always right-sided (only large values speak against H_0)
- The distribution of F is known under H_0 : the Fisher-distribution with $(k-1)$ -numerator- and $(n-k)$ -denominator- degrees of freedom, short


$$F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k) \quad (\text{R knows it well, } \text{rf}(\dots), \text{ pf}(\dots) \text{ etc.})$$

For more details on the $\mathcal{F}(k-1, n-k)$ -distribution and the derivation of the distribution of F see e.g.,
Messer, M. and Schneider, G. *Statistik: Theorie und Praxis im Dialog*, Springer Berlin

F-test



Under H_0 : $F \sim F(k-1, n-k)$




- Null hypothesis $H_0 : \mu_1 = \dots = \mu_k$
- F-statistic


$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$

F-test



Under H_0 : $F \sim F(k-1, n-k)$




- Null hypothesis $H_0 : \mu_1 = \dots = \mu_k$
- F-statistic


$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{ij} - \bar{X}_{i\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$

F-test



Under H_0 : $F \sim F(k-1, n-k)$




- Null hypothesis $H_0 : \mu_1 = \dots = \mu_k$
- F-statistic


$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{ij} - \bar{X}_{i\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$
 - means: $\bar{x}_{1\cdot} \approx 50.8$, $\bar{x}_{2\cdot} \approx 61.3$, $\bar{x}_{3\cdot} \approx 55.5$, $\bar{x}_{4\cdot} \approx 50.0$ and $\bar{x} \approx 53.3$

F-test



Under H_0 : $F \sim F(k-1, n-k)$





- Null hypothesis $H_0 : \mu_1 = \dots = \mu_k$
- F -statistic

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$
 - means: $\bar{x}_{1,\cdot} \approx 50.8$, $\bar{x}_{2,\cdot} \approx 61.3$, $\bar{x}_{3,\cdot} \approx 55.5$, $\bar{x}_{4,\cdot} \approx 50.0$ and $\bar{x} \approx 53.3$
 - thus: $f \approx 7.0$

F-test





- Null hypothesis $H_0: \mu_1 = \dots = \mu_k$
- F-statistic

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{ij} - \bar{X}_{i\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$
 - means: $\bar{x}_{1\cdot} \approx 50.8$, $\bar{x}_{2\cdot} \approx 61.3$, $\bar{x}_{3\cdot} \approx 55.5$, $\bar{x}_{4\cdot} \approx 50.0$ and $\bar{x} \approx 53.3$
 - thus: $f \approx 7.0$
- For $\alpha = 5\%$ it is $q_{1-\alpha} \approx 2.69$ (95%-quantile of $\mathcal{F}(3, 98)$), thus $R \approx [2.69, \infty)$

F-test





- Null hypothesis $H_0 : \mu_1 = \dots = \mu_k$
- F-statistic

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{ij} - \bar{X}_{i,\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$
 - means: $\bar{x}_{1,\cdot} \approx 50.8$, $\bar{x}_{2,\cdot} \approx 61.3$, $\bar{x}_{3,\cdot} \approx 55.5$, $\bar{x}_{4,\cdot} \approx 50.0$ and $\bar{x} \approx 53.3$
 - thus: $f \approx 7.0$
- For $\alpha = 5\%$ it is $q_{1-\alpha} \approx 2.69$ (95%-quantile of $\mathcal{F}(3, 98)$), thus $R \approx [2.69, \infty)$
- $f \in R$, and hence reject H_0 on the 5%-level

F-test





- Null hypothesis $H_0: \mu_1 = \dots = \mu_k$
- F -statistic

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{ij} - \bar{X}_{i\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$
 - means: $\bar{x}_{1\cdot} \approx 50.8$, $\bar{x}_{2\cdot} \approx 61.3$, $\bar{x}_{3\cdot} \approx 55.5$, $\bar{x}_{4\cdot} \approx 50.0$ and $\bar{x} \approx 53.3$
 - thus: $f \approx 7.0$
- For $\alpha = 5\%$ it is $q_{1-\alpha} \approx 2.69$ (95%-quantile of $\mathcal{F}(3, 98)$), thus $R \approx [2.69, \infty)$
- $f \in R$, and hence reject H_0 on the 5%-level
- $p = \mathbb{P}_{H_0}(F \geq f) \approx 0.00025 = 1/4000$. Only in about one of 4000 cases we observe a discrepancy, which is at least as extreme as in the data, if the null hypothesis holds true.

F-test



- Null hypothesis $H_0: \mu_1 = \dots = \mu_k$
- F-statistic


$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{ij} - \bar{X}_{i\cdot})^2}$$

- $F \stackrel{H_0}{\sim} \mathcal{F}(k-1, n-k)$
- Evaluation of the data yields f
 - $k = 4$ groups, $n_1 = 25$, $n_2 = 16$, $n_3 = 25$, $n_4 = 36$, thus $n = 102$
 - means: $\bar{x}_{1\cdot} \approx 50.8$, $\bar{x}_{2\cdot} \approx 61.3$, $\bar{x}_{3\cdot} \approx 55.5$, $\bar{x}_{4\cdot} \approx 50.0$ and $\bar{x} \approx 53.3$
 - thus: $f \approx 7.0$
- For $\alpha = 5\%$ it is $q_{1-\alpha} \approx 2.69$ (95%-quantile of $\mathcal{F}(3, 98)$), thus $R \approx [2.69, \infty)$
- $f \in R$, and hence reject H_0 on the 5%-level
- $p = \mathbb{P}_{H_0}(F \geq f) \approx 0.00025 = 1/4000$. Only in about one of 4000 cases we observe a discrepancy, which is at least as extreme as in the data, if the null hypothesis holds true.
- Result plausible, given the **variability (sem)** of the **means**

ANOVA - Summary

Model:


- Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$



ANOVA - Summary

Model:

- Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus, k groups and $n = \sum_{i=1}^k n_i$ observations in total




ANOVA - Summary

Model:


- Let $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$ be independent RVs and for $i = 1, \dots, k$ let $X_{i,j} \sim N(\mu_i, \sigma^2)$ for $j = 1, \dots, n_i$, with $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus, k groups and $n = \sum_{i=1}^k n_i$ observations in total

Under $H_0 : \mu_1 = \dots = \mu_k$ it holds

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} \sim \mathcal{F}(k-1, n-k)$$



Under $H_0 : F \sim \mathcal{F}(k-1, n-k)$



ANOVA in R

```
# Sizes of groups as vector (k groups)
ni      <- c(n1,n2,...,nk)
# Data as vector, all from group 1, then all from group 2 etc..
x       <- c(...)
# Groups as factor variable
gr      <- factor(rep(1:k,ni))
# Perform ANOVA (F-test)
anova(aov(x~gr))
```

Output

Analysis of Variance Table

Response: x

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
gr	3	1709.8	569.92	7.0332	0.000248 ***
Residuals	98	7941.2	81.03		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

- 1. column: degrees of freedom (first row numerator, second row denominator)
 - 3. column: numerator and denominator of the f-statistic
 - 4. column: f-statistic
 - 5. column: p -value
- (2.column = 1.column · 3.column)

In practice

Procedure in practice often:

- First apply the ANOVA...

In practice

Procedure in practice often:

- First apply the ANOVA...
- ...if the null hypothesis is rejected, then perform post-hoc t -tests (and correct the p -values e.g., using Bonferroni correction)

In practice

Procedure in practice often:

- First apply the ANOVA...
- ...if the null hypothesis is rejected, then perform post-hoc t -tests (and correct the p -values e.g., using Bonferroni correction)

Rethink of whether the data comply with the model assumptions of the ANOVA:

- Independence and own expectation per group
→ makes sense due to the practical setup


In practice

Procedure in practice often:

- First apply the ANOVA...
- ...if the null hypothesis is rejected, then perform post-hoc t -tests (and correct the p -values e.g., using Bonferroni correction)

Rethink of whether the data comply with the model assumptions of the ANOVA:

- Independence and own expectation per group
→ makes sense due to the practical setup
- normal distribution and constant variance (between the groups)
→ plausible when looking at the data



Thank you!