


Probability

Conditional probability and the Bayes theorem



Terminology

- **Random experiment** ... a repeatable procedure with well-defined possible outcomes

Terminology

- **Random experiment** ... a repeatable procedure with well-defined possible outcomes
- **Sample space Ω** ... the set of all possible outcomes
 - **Discrete sample space** - countable (listable) and it can be finite or infinite
 - Example: $\{H, T\}$, $\{1, 3, 5, \dots\}$ are discrete sample spaces
The interval $0 \leq x \leq 1$ is not discrete, but continuous
 - Describe (list) all possible outcomes
 - Describe the probability of the outcomes

... **Probability tables**

Terminology

- **Random experiment** ... a repeatable procedure with well-defined possible outcomes
- **Sample space Ω** ... the set of all possible outcomes
 - **Discrete sample space** - countable (listable) and it can be finite or infinite
 - Example: $\{H, T\}$, $\{1, 3, 5, \dots\}$ are discrete sample spaces
The interval $0 \leq x \leq 1$ is not discrete, but continuous
 - Describe (list) all possible outcomes
 - Describe the probability of the outcomes
- **Event A** ... a subset of the sample space
 - Events are sets that can be described in: words, notation, with Venn diagrams

... **Probability tables**

Terminology

- **Random experiment** ... a repeatable procedure with well-defined possible outcomes
 - **Sample space Ω** ... the set of all possible outcomes
 - **Discrete sample space** - countable (listable) and it can be finite or infinite
 - Example: $\{H, T\}, \{1, 3, 5, \dots\}$ are discrete sample spaces
The interval $0 \leq x \leq 1$ is not discrete, but continuous
 - Describe (list) all possible outcomes
 - Describe the probability of the outcomes
- ... **Probability tables**
- **Event A** ... a subset of the sample space
 - Events are sets that can be described in: words, notation, with Venn diagrams
 - **Probability mass function** ... a function that gives the probability of each outcome
 $P: \mathcal{P}(\Omega) \rightarrow [0, 1]$

Terminology

- **Random experiment** ... a repeatable procedure with well-defined possible outcomes
- **Sample space Ω** ... the set of all possible outcomes
 - **Discrete sample space** - countable (listable) and it can be finite or infinite
 - Example: $\{H, T\}, \{1, 3, 5, \dots\}$ are discrete sample spaces
The interval $0 \leq x \leq 1$ is not discrete, but continuous
 - Describe (list) all possible outcomes
 - Describe the probability of the outcomes

... **Probability tables**

- **Event A** ...a subset of the sample space
 - Events are sets that can be described in: words, notation, with Venn diagrams
- **Probability mass function** ...a function that gives the probability of each outcome
 $P: \mathcal{P}(\Omega) \rightarrow [0, 1]$

Axioms: $P(\Omega) = 1$; $P(A) \geq 0$ for all A ; and
 $P(A \cup B) = P(A) + P(B)$ for all disjoint A, B

Terminology

- **Random experiment** ... a repeatable procedure with well-defined possible outcomes
 - **Sample space Ω** ... the set of all possible outcomes
 - **Discrete sample space** - countable (listable) and it can be finite or infinite
 - Example: $\{H, T\}, \{1, 3, 5, \dots\}$ are discrete sample spaces
The interval $0 \leq x \leq 1$ is not discrete, but continuous
 - Describe (list) all possible outcomes
 - Describe the probability of the outcomes
- ... **Probability tables**
- **Event A** ...a subset of the sample space
 - Events are sets that can be described in: words, notation, with Venn diagrams
 - **Probability mass function** ...a function that gives the probability of each outcome
 $P: \mathcal{P}(\Omega) \rightarrow [0, 1]$
Axioms: $P(\Omega) = 1$; $P(A) \geq 0$ for all A ; and
 $P(A \cup B) = P(A) + P(B)$ for all disjoint A, B
 - $(\Omega, \mathcal{P}(\Omega), P)$ is called a **probability space**

Discrete probability space

- Let $\Omega = \{\omega_1, \omega_2, \dots\}$ countable. Let p_i be the probability $p_i = P(\omega_i)$.

Discrete probability space

- Let $\Omega = \{\omega_1, \omega_2, \dots\}$ countable. Let p_i be the probability $p_i = P(\omega_i)$.
 - The probability (mass function) P is **uniquely** determined by the probabilities of the elementary events

Discrete probability space

- Let $\Omega = \{\omega_1, \omega_2, \dots\}$ countable. Let p_i be the probability $p_i = P(\omega_i)$.
- The probability (mass function) P is uniquely determined by the probabilities of the elementary events
- Probability table

elementary events	ω_i	ω_1	ω_2	\dots	ω_n	\dots
probability	p_i	p_1	p_2	\dots	p_n	\dots

Discrete probability space

- Let $\Omega = \{\omega_1, \omega_2, \dots\}$ countable. Let p_i be the probability $p_i = P(\omega_i)$.
 - The probability (mass function) P is **uniquely** determined by the probabilities of the elementary events
 - **Probability table**

elementary events	ω_i	ω_1	ω_2	\dots	ω_n	\dots
probability	p_i	p_1	p_2	\dots	p_n	\dots

$$0 \leq p_i \leq 1 \quad \text{and} \quad \sum_i p_i = 1$$

Discrete probability space

- Let $\Omega = \{\omega_1, \omega_2, \dots\}$ countable. Let p_i be the probability $p_i = P(\omega_i)$.
- The probability (mass function) P is **uniquely** determined by the probabilities of the elementary events

- **Probability table**

elementary events	ω_i	ω_1	ω_2	\dots	ω_n	\dots
probability	p_i	p_1	p_2	\dots	p_n	\dots

$$0 \leq p_i \leq 1 \quad \text{and} \quad \sum_i p_i = 1$$

- Probability of an event A

$$P(A) = \sum_{\omega_i \in A} p_i$$

Simple examples

- (1) Let us toss a fair coin twice.

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

- **Event A:** head appeared once

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

- **Event A:** head appeared once

$$A = \{HT, TH\} \quad \text{and} \quad P(A) = \frac{1}{2}$$

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

- **Event A:** head appeared once

$$A = \{HT, TH\} \quad \text{and} \quad P(A) = \frac{1}{2}$$

- **Event B:** head appeared at least once

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

- **Event A:** head appeared once

$$A = \{HT, TH\} \quad \text{and} \quad P(A) = \frac{1}{2}$$

- **Event B:** head appeared at least once

$$B = \{HH, HT, TH\} \quad \text{and} \quad P(B) = \frac{3}{4}$$

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

- **Event A:** head appeared once

$$A = \{HT, TH\} \quad \text{and} \quad P(A) = \frac{1}{2}$$

- **Event B:** head appeared at least once

$$B = \{HH, HT, TH\} \quad \text{and} \quad P(B) = \frac{3}{4}$$

- **Event C:** head appeared three times

Simple examples

(1) Let us toss a fair coin twice.

- **Random experiment:** toss a fair coin twice, list the outcomes
- **Sample space:** $\Omega = \{HH, HT, TH, TT\}$ and $|\Omega| = 4$
- **Probability:** each outcome is equally likely with probability $\frac{1}{4}$

ω_i	HH	HT	TH	TT	
p_i	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	1

- **Event A:** head appeared once

$$A = \{HT, TH\} \quad \text{and} \quad P(A) = \frac{1}{2}$$

- **Event B:** head appeared at least once

$$B = \{HH, HT, TH\} \quad \text{and} \quad P(B) = \frac{3}{4}$$

- **Event C:** head appeared three times

$$C = \emptyset \quad \text{and} \quad P(C) = 0$$

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

- **Probability:**

ω_i	H	TH	TTH	TTTH	TTTTH	...
p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

• Probability:	ω_i	H	TH	TTH	TTTH	TTTTH	...
	p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Is the sum of all probabilities equal 1?

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

- **Probability:**

ω_i	H	TH	TTH	TTTH	TTTTH	...
p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Is the sum of all probabilities equal 1?

We use geometric series:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots = \frac{1}{2} \cdot (1 + \frac{1}{2} + \frac{1}{2^2} + \dots) = \frac{1}{2} \cdot \frac{1}{1-\frac{1}{2}} = 1$$

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

- **Probability:**

ω_i	H	TH	TTH	TTTH	TTTTH	...
p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Is the sum of all probabilities equal 1?

We use geometric series:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots = \frac{1}{2} \cdot (1 + \frac{1}{2} + \frac{1}{2^2} + \dots) = \frac{1}{2} \cdot \frac{1}{1-\frac{1}{2}} = 1$$

- **Event A:** head was obtained in a maximum of 4 tosses

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

- **Probability:**

ω_i	H	TH	TTH	TTTH	TTTTH	...
p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Is the sum of all probabilities equal 1?

We use geometric series:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots = \frac{1}{2} \cdot (1 + \frac{1}{2} + \frac{1}{2^2} + \dots) = \frac{1}{2} \cdot \frac{1}{1-\frac{1}{2}} = 1$$

- **Event A:** head was obtained in a maximum of 4 tosses

$$A = \{H, TH, TTH, TTTH\} \quad \text{and} \quad P(A) = \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} = 0.9375$$

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

- **Probability:**

ω_i	H	TH	TTH	TTTH	TTTTH	...
p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Is the sum of all probabilities equal 1?

We use geometric series:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots = \frac{1}{2} \cdot (1 + \frac{1}{2} + \frac{1}{2^2} + \dots) = \frac{1}{2} \cdot \frac{1}{1-\frac{1}{2}} = 1$$

- **Event A:** head was obtained in a maximum of 4 tosses

$$A = \{H, TH, TTH, TTTH\} \quad \text{and} \quad P(A) = \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} = 0.9375$$

- **Event B:** head appears in at least 3 throws

Simple examples

(2) We toss a fair coin (infinite sample space)

- **Random experiment:** We toss a fair coin until the first heads.
- **Sample space:** $\Omega = \{H, TH, TTH, TTTH, \dots\}$

- **Probability:**

ω_i	H	TH	TTH	TTTH	TTTTH	...
p_i	$\frac{1}{2}$	$\frac{1}{2^2}$	$\frac{1}{2^3}$	$\frac{1}{2^4}$	$\frac{1}{2^5}$...

Is the sum of all probabilities equal 1?

We use geometric series:

$$\frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \dots = \frac{1}{2} \cdot (1 + \frac{1}{2} + \frac{1}{2^2} + \dots) = \frac{1}{2} \cdot \frac{1}{1-\frac{1}{2}} = 1$$

- **Event A:** head was obtained in a maximum of 4 tosses

$$A = \{H, TH, TTH, TTTH\} \quad \text{and} \quad P(A) = \frac{1}{2} + \frac{1}{2^2} + \frac{1}{2^3} + \frac{1}{2^4} = 0.9375$$

- **Event B:** head appears in at least 3 throws

$$B = \Omega \setminus \{H, TH\} \quad \text{and} \quad P(B) = 1 - (\frac{1}{2} + \frac{1}{2^2}) = 0.25$$

Simple examples

- (3) Taxis (an infinite discrete sample space)

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.
- **Probability:** this is often modelled using the probability mass function

$$p_k = P(k) = e^{-\lambda} \cdot \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

known as the **Poisson distribution**, where λ is the average number of taxis.

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.
- **Probability:** this is often modelled using the probability mass function

$$p_k = P(k) = e^{-\lambda} \cdot \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

known as the **Poisson distribution**, where λ is the average number of taxis.

- We can put this in a table:

ω_i	0	1	...	n	...
p_i	$e^{-\lambda}$	$e^{-\lambda} \cdot \lambda$...	$e^{-\lambda} \cdot \frac{\lambda^n}{n!}$...

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.
- **Probability:** this is often modelled using the probability mass function

$$p_k = P(k) = e^{-\lambda} \cdot \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

known as the **Poisson distribution**, where λ is the average number of taxis.

- We can put this in a table:

ω_i	0	1	...	n	...
p_i	$e^{-\lambda}$	$e^{-\lambda} \cdot \lambda$...	$e^{-\lambda} \cdot \frac{\lambda^n}{n!}$...

Is the sum of the probabilities of all possible outcomes 1?

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.
- **Probability:** this is often modelled using the probability mass function

$$p_k = P(k) = e^{-\lambda} \cdot \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

known as the **Poisson distribution**, where λ is the average number of taxis.

- We can put this in a table:

ω_i	0	1	...	n	...
p_i	$e^{-\lambda}$	$e^{-\lambda} \cdot \lambda$...	$e^{-\lambda} \cdot \frac{\lambda^n}{n!}$...

Is the sum of the probabilities of all possible outcomes 1? Yes!

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.
- **Probability:** this is often modelled using the probability mass function

$$p_k = P(k) = e^{-\lambda} \cdot \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

known as the **Poisson distribution**, where λ is the average number of taxis.

- We can put this in a table:

ω_i	0	1	...	n	...
p_i	$e^{-\lambda}$	$e^{-\lambda} \cdot \lambda$...	$e^{-\lambda} \cdot \frac{\lambda^n}{n!}$...

Is the sum of the probabilities of all possible outcomes 1? Yes!

Using the Taylor series $e^\lambda = \sum_{k=0}^{\infty} \frac{\lambda^k}{k!}$, we obtain

$$\sum_{k=0}^{\infty} p_k = e^{-\lambda} \cdot \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} = e^{-\lambda} \cdot e^\lambda = 1$$

Simple examples

(3) Taxis (an infinite discrete sample space)

- **Random experiment:** count the number of taxis that pass through Wiedner Hauptstrasse during our class
- **Sample space:** $\Omega = \{0, 1, 2, \dots\}$.
- **Probability:** this is often modelled using the probability mass function

$$p_k = P(k) = e^{-\lambda} \cdot \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

known as the **Poisson distribution**, where λ is the average number of taxis.

- We can put this in a table:

ω_i	0	1	...	n	...
p_i	$e^{-\lambda}$	$e^{-\lambda} \cdot \lambda$...	$e^{-\lambda} \cdot \frac{\lambda^n}{n!}$...

Is the sum of the probabilities of all possible outcomes 1? Yes!

Using the Taylor series $e^\lambda = \sum_{k=0}^{\infty} \frac{\lambda^k}{k!}$, we obtain

$$\sum_{k=0}^{\infty} p_k = e^{-\lambda} \cdot \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} = e^{-\lambda} \cdot e^\lambda = 1$$

HW What is the probability that 2 taxis will pass by? 5? at least 8?

Simple examples

- (4) Two dice (choice of sample space)

Simple examples

(4) Two dice (choice of sample space)

- Random experiment (a): roll two dice, note the pair of numbers showing on the dice (first dice, second dice)

Simple examples

(4) Two dice (choice of sample space)

- **Random experiment (a)**: roll two dice, note the **pair of numbers** showing on the dice (first dice, second dice)
- **Sample space**: the product of the result sets for each die

$$\Omega = \{(1,1), (2,1), \dots, (6,5), (6,6)\}, \quad |\Omega| = 36$$

Simple examples

(4) Two dice (choice of sample space)

- **Random experiment (a)**: roll two dice, note the **pair of numbers** showing on the dice (first dice, second dice)
- **Sample space**: the product of the result sets for each die

$$\Omega = \{(1, 1), (2, 1), \dots, (6, 5), (6, 6)\}, \quad |\Omega| = 36$$

- The **probability** of each outcome is $\frac{1}{36}$.

Simple examples

(4) Two dice (choice of sample space)

- **Random experiment (a):** roll two dice, note the **pair of numbers** showing on the dice (first die, second die)
- **Sample space:** the product of the result sets for each die

$$\Omega = \{(1, 1), (2, 1), \dots, (6, 5), (6, 6)\}, \quad |\Omega| = 36$$

- The **probability** of each outcome is $\frac{1}{36}$.

		die 2					
		1	2	3	4	5	6
die 1	1	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	2	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	3	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	4	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	5	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	6	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$

two dimensional
probability table (Table 1)

Simple examples

(4) Two dice (choice of sample space)

- Random experiment (a): roll two dice, note the pair of numbers showing on the dice (first die, second die)

- Sample space: the product of the result sets for each die

$$\Omega = \{(1, 1), (2, 1), \dots, (6, 5), (6, 6)\}, \quad |\Omega| = 36$$

- The probability of each outcome is $\frac{1}{36}$.

		die 2					
		1	2	3	4	5	6
die 1	1	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	2	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	3	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	4	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	5	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	6	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$

two dimensional
probability table (Table 1)

- A: one gets the same number on both dice

Simple examples

(4) Two dice (choice of sample space)

- Random experiment (a): roll two dice, note the pair of numbers showing on the dice (first die, second die)

- Sample space: the product of the result sets for each die

$$\Omega = \{(1,1), (2,1), \dots, (6,5), (6,6)\}, \quad |\Omega| = 36$$

- The probability of each outcome is $\frac{1}{36}$.

		die 2					
		1	2	3	4	5	6
die 1	1	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	2	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	3	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	4	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	5	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$
	6	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$	$\frac{1}{36}$

two dimensional
probability table (Table 1)

- A: one gets the same number on both dice

$$A = \{(1,1), (2,2), (3,3), (4,4), (5,5), (6,6)\} \quad \text{and} \quad P(A) = \frac{6}{36} = \frac{1}{6}$$

Simple examples

(5) Two dice (choice of sample space)

Simple examples

(5) Two dice (choice of sample space)

- Random experiment (b): roll two dice and note the sum of the numbers on the dice

Simple examples

(5) Two dice (choice of sample space)

- Random experiment (b): roll two dice and note the sum of the numbers on the dice
- Sample space: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$, $|\Omega| = 11$

Simple examples

(5) Two dice (choice of sample space)

- **Random experiment (b)**: roll two dice and note the **sum** of the numbers on the dice
- **Sample space**: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$, $|\Omega| = 11$
- **probability**: these 11 outcomes are *not* all equally likely.

ω_i	2	3	4	5	6	7	8	9	10	11	12	
p_i	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$	1

Table 2

Simple examples

(5) Two dice (choice of sample space)

- **Random experiment (b)**: roll two dice and note the **sum** of the numbers on the dice
- **Sample space**: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$, $|\Omega| = 11$
- **probability**: these 11 outcomes are *not* all equally likely.

ω_i	2	3	4	5	6	7	8	9	10	11	12	
p_i	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$	1

Table 2

- **A**: the sum is smaller than 6

Simple examples

(5) Two dice (choice of sample space)

- **Random experiment (b)**: roll two dice and note the **sum** of the numbers on the dice
- **Sample space**: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$, $|\Omega| = 11$
- **probability**: these 11 outcomes are *not* all equally likely.

ω_i	2	3	4	5	6	7	8	9	10	11	12	
p_i	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$	1

Table 2

- **A**: the sum is smaller than 6
 - $A = \{2, 3, 4, 5\}$ and $P(A) = \frac{1}{36} + \frac{2}{36} + \frac{3}{36} + \frac{3}{36} = \frac{5}{18}$

Simple examples

(5) Two dice (choice of sample space)

- **Random experiment (b)**: roll two dice and note the **sum** of the numbers on the dice
- **Sample space**: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$, $|\Omega| = 11$
- **probability**: these 11 outcomes are *not* all equally likely.

ω_i	2	3	4	5	6	7	8	9	10	11	12	
p_i	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$	1

Table 2

- **A**: the sum is smaller than 6
 - $A = \{2, 3, 4, 5\}$ and $P(A) = \frac{1}{36} + \frac{2}{36} + \frac{3}{36} + \frac{3}{36} = \frac{5}{18}$

HW

- **B**: The total is even
- **C**: the sum is at most 10
- **D**: the sum is 1
- Are A and C disjoint? Compute $A \cap B$, $B \cup C$ and $B^c \cap D$.

Simple examples

(5) Two dice (choice of sample space)

- **Random experiment (b)**: roll two dice and note the **sum** of the numbers on the dice
- **Sample space**: $\Omega = \{2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12\}$, $|\Omega| = 11$
- **probability**: these 11 outcomes are *not* all equally likely.

ω_i	2	3	4	5	6	7	8	9	10	11	12	
p_i	$\frac{1}{36}$	$\frac{2}{36}$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{5}{36}$	$\frac{6}{36}$	$\frac{5}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	$\frac{2}{36}$	$\frac{1}{36}$	1

Table 2

- **A**: the sum is smaller than 6
 - $A = \{2, 3, 4, 5\}$ and $P(A) = \frac{1}{36} + \frac{2}{36} + \frac{3}{36} + \frac{3}{36} = \frac{5}{18}$

HW

- **B**: The total is even
- **C**: the sum is at most 10
- **D**: the sum is 1
- Are A and C disjoint? Compute $A \cap B$, $B \cup C$ and $B^c \cap D$.
- **Question**: What is the relationship between the two probability tables above?

Simple examples

HW The statistical experiment

- **Experiment:** random selection of k objects from a set of n distinguishable objects $\mathcal{M} = \{1, 2, \dots, n\}$, $k \leq n$, assuming an object that has been selected cannot be selected a second time.

Describe the result sets and determine their size when selected a second time.

- the order of selection does not matter
- the order of selection is important.

Simple examples

- (6) Measure the lifetime of a transistor
(Real probability space)

Simple examples

- (6) Measure the lifetime of a transistor
(Real probability space)

... continuous space

Simple examples

- (6) Measure the lifetime of a transistor
(Real probability space)

... continuous space

- **Random experiment:** measure the lifetime (in operating hours) of a transistor and report the result

Simple examples

- (6) Measure the lifetime of a transistor
(Real probability space)

... continuous space

- **Random experiment:** measure the lifetime (in operating hours) of a transistor and report the result
- **Sample space:** $\Omega = [0, \infty)$, i.e. we can obtain any nonnegative real value

Simple examples

- (6) Measure the lifetime of a transistor
(Real probability space)

... continuous space

- **Random experiment:** measure the lifetime (in operating hours) of a transistor and report the result
- **Sample space:** $\Omega = [0, \infty)$, i.e. we can obtain any nonnegative real value
- **Probability:** Since there is a continuum of possible outcomes, we have to use a probability density function.

Simple examples

- (6) Measure the lifetime of a transistor
(Real probability space)

... continuous space

- **Random experiment:** measure the lifetime (in operating hours) of a transistor and report the result
- **Sample space:** $\Omega = [0, \infty)$, i.e. we can obtain any nonnegative real value
- **Probability:** Since there is a continuum of possible outcomes, we have to use a probability density function.

We will talk about it later.

Some rules of probability

- Let A , B and C be events contained in Ω .

Some rules of probability

- Let A , B and C be events contained in Ω .

(1) $P(A^c) = 1 - P(A)$

Some rules of probability

- Let A , B and C be events contained in Ω .

(1) $P(A^c) = 1 - P(A)$

(2) $P(B \cup C) = P(B) + P(C) - P(B \cap C)$ for all B, C

Some rules of probability

- Let A , B and C be events contained in Ω .

(1) $P(A^c) = 1 - P(A)$

(2) $P(B \cup C) = P(B) + P(C) - P(B \cap C)$ for all B, C

(3) $A \subseteq B \implies P(A) \leq P(B)$

Some rules of probability

- Let A , B and C be events contained in Ω .

(1) $P(A^c) = 1 - P(A)$

(2) $P(B \cup C) = P(B) + P(C) - P(B \cap C)$ for all B, C

(3) $A \subseteq B \implies P(A) \leq P(B)$

We can visualize these rules using [Venn diagrams](#).

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?
 - If A is an event of being an odd random number, then A^c is an event of being an even random number. Then, $P(A^c) = 1 - P(A) = 0.35$.

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?
 - If A is an event of being an odd random number, then A^c is an event of being an even random number. Then, $P(A^c) = 1 - P(A) = 0.35$.
 - (b) Consider two events, A is an integer divisible by two and B is odd and less than 10. Suppose $P(A) = 0.6$ and $P(B) = 0.25$. What is $A \cap B$? What is the probability $P(A \cup B)$?

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?
 - If A is an event of being an odd random number, then A^c is an event of being an even random number. Then, $P(A^c) = 1 - P(A) = 0.35$.
 - (b) Consider two events, A is an integer divisible by two and B is odd and less than 10. Suppose $P(A) = 0.6$ and $P(B) = 0.25$. What is $A \cap B$? What is the probability $P(A \cup B)$?
 - A and B are **disjoint**, since all numbers in A are even and all numbers in B are odd, i.e. $A \cap B = \emptyset$.

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?
 - If A is an event of being an odd random number, then A^c is an event of being an even random number. Then, $P(A^c) = 1 - P(A) = 0.35$.
 - (b) Consider two events, A is an integer divisible by two and B is odd and less than 10. Suppose $P(A) = 0.6$ and $P(B) = 0.25$. What is $A \cap B$? What is the probability $P(A \cup B)$?
 - A and B are **disjoint**, since all numbers in A are even and all numbers in B are odd, i.e. $A \cap B = \emptyset$.
 - $P(A \cup B) = P(A) + P(B) = 0.85$

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?
 - If A is an event of being an odd random number, then A^c is an event of being an even random number. Then, $P(A^c) = 1 - P(A) = 0.35$.
 - (b) Consider two events, A is an integer divisible by two and B is odd and less than 10. Suppose $P(A) = 0.6$ and $P(B) = 0.25$. What is $A \cap B$? What is the probability $P(A \cup B)$?
 - A and B are **disjoint**, since all numbers in A are even and all numbers in B are odd, i.e. $A \cap B = \emptyset$.
 - $P(A \cup B) = P(A) + P(B) = 0.85$
 - (c) Consider three events, A is a random number and is a multiple of 2, B is a multiple of 3 and C is a multiple of 6. Suppose $P(A) = 0.6$ and $P(B) = 0.3$ and $P(C) = 0.2$. What is the probability of the event A or B ?

Examples

- For the following examples suppose we have an experiment that produces a random integer between 1 and 20. The probabilities are not necessarily the same for each outcome.
 - (a) If the probability of an odd number is 0.65 what is the probability of an even number?
 - If A is an event of being an odd random number, then A^c is an event of being an even random number. Then, $P(A^c) = 1 - P(A) = 0.35$.
 - (b) Consider two events, A is an integer divisible by two and B is odd and less than 10. Suppose $P(A) = 0.6$ and $P(B) = 0.25$. What is $A \cap B$? What is the probability $P(A \cup B)$?
 - A and B are **disjoint**, since all numbers in A are even and all numbers in B are odd, i.e. $A \cap B = \emptyset$.
 - $P(A \cup B) = P(A) + P(B) = 0.85$
 - (c) Consider three events, A is a random number and is a multiple of 2, B is a multiple of 3 and C is a multiple of 6. Suppose $P(A) = 0.6$ and $P(B) = 0.3$ and $P(C) = 0.2$. What is the probability of the event A or B ?
 - Since an integer is divisible by 6 if and only if it is divisible by both 2 and 3, we have $C = A \cap B$. Then,
 $P(A \text{ or } B) = P(A \cup B) = P(A) + P(B) - P(C) = 0.6 + 0.3 - 0.2 = 0.7$.

Examples

HW If $P(A^c) = \frac{1}{4}$ and $P(B) = \frac{1}{3}$, can A and B be disjoint?

Justify your answer.

HW The 24h express delivery has increased significantly in the last ten years. Customer service has consistently been shown to be the greatest influence on a company's success. A study was conducted to examine customer satisfaction with an express delivery service. In addition to their satisfaction, the customers were asked how often they had used the overnight shipping option. The results are shown below

Frequency of Use	Satisfaction level			Total
	High	Medium	Low	
< 2 per month	250	140	10	400
2-5 per month	140	55	5	200
> 5 per month	70	25	5	100
Total	460	220	20	700

- What is the probability that a randomly selected person will use the company two to five times a month or have moderate satisfaction?

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω

In total $|\Omega| = \binom{52}{5}$ poker hands

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) **A:** we got exactly one pair

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) A : we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) A : we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

- 13 ... number of ways to select the pair

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♡, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) A : we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

- 13 ... number of ways to select the pair
- $\binom{4}{2}$... number of ways to choose the two cards from the pair

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♡, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) *A*: we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

- 13 ... number of ways to select the pair
- $\binom{4}{2}$... number of ways to choose the two cards from the pair
- $\binom{12}{3}$... number of ways to specify the other three cards

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) *A*: we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

- 13 ... number of ways to select the pair
- $\binom{4}{2}$... number of ways to choose the two cards from the pair
- $\binom{12}{3}$... number of ways to specify the other three cards
- 4^3 ... number of ways to choose the other three cards from these values

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) A: we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

- 13 ... number of ways to select the pair
- $\binom{4}{2}$... number of ways to choose the two cards from the pair
- $\binom{12}{3}$... number of ways to specify the other three cards
- 4^3 ... number of ways to choose the other three cards from these values

$$P(\text{exactly one pair}) = \frac{13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3}{\binom{52}{5}} = \frac{1098240}{2598960} \approx 42.3\%$$

Another example

(1) Playing poker

52 cards: 13 ranks 2, ..., 9, 10, J, Q, K, A and 4 suits ♥, ♠, ♦, ♣

Poker hands consist of 5 cards

- **Random experiment:** playing poker with cards
- **Sample space:** Ω contains all poker hands ω
In total $|\Omega| = \binom{52}{5}$ poker hands
- **Probability:** all $\omega \in \Omega$ are equally probable

(a) *A*: we got exactly one pair

The number of hands with exactly one pair is

$$13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3 = 1098240$$

- 13 ... number of ways to select the pair
- $\binom{4}{2}$... number of ways to choose the two cards from the pair
- $\binom{12}{3}$... number of ways to specify the other three cards
- 4^3 ... number of ways to choose the other three cards from these values

$$P(\text{exactly one pair}) = \frac{13 \cdot \binom{4}{2} \cdot \binom{12}{3} \cdot 4^3}{\binom{52}{5}} = \frac{1098240}{2598960} \approx 42.3\%$$

... note, a full house is not a hand with exactly one pair

Another example

(b) B : we get four kings

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings?

Another example

(b) B : we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
- If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
- If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

- We can also calculate the probability through an “updating” argument, by calculating conditional probabilities.

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
 - If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

- We can also calculate the probability through an “updating” argument, by calculating conditional probabilities.
 - The probability that the first card is a king is $\frac{4}{52}$.

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
- If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

- We can also calculate the probability through an “updating” argument, by calculating conditional probabilities.
- The probability that the first card is a king is $\frac{4}{52}$. Assuming the first card is a king, the probability that the second card is a king is $\frac{3}{51}$.

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
- If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

- We can also calculate the probability through an “updating” argument, by calculating conditional probabilities.
- The probability that the first card is a king is $\frac{4}{52}$. Assuming the first card is a king, the probability that the second card is a king is $\frac{3}{51}$. If we continue this argument, we get that the probability is four kings

$$P(\text{four kings}) = \frac{4}{52} \cdot \frac{3}{51} \cdot \frac{2}{50} \cdot \frac{1}{49} \cdot \frac{48}{48} \cdot \binom{5}{1} = \frac{1}{54145}.$$

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
- If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

- We can also calculate the probability through an “updating” argument, by calculating conditional probabilities.
- The probability that the first card is a king is $\frac{4}{52}$. Assuming the first card is a king, the probability that the second card is a king is $\frac{3}{51}$. If we continue this argument, we get that the probability is four kings

$$P(\text{four kings}) = \frac{4}{52} \cdot \frac{3}{51} \cdot \frac{2}{50} \cdot \frac{1}{49} \cdot \frac{48}{48} \cdot \binom{5}{1} = \frac{1}{54145}.$$

Another example

(b) *B*: we get four kings

- How many different hands are there with four kings? What is the probability of having four kings?
- If we specify that four of the cards are kings, there are 48 different ways to determine the fifth card. Thus,

$$P(\text{four kings}) = \frac{48}{\binom{52}{5}} = \frac{1}{54145} < \frac{1}{50000}.$$

- We can also calculate the probability through an “updating” argument, by calculating conditional probabilities.
- The probability that the first card is a king is $\frac{4}{52}$. Assuming the first card is a king, the probability that the second card is a king is $\frac{3}{51}$. If we continue this argument, we get that the probability is four kings

$$P(\text{four kings}) = \frac{4}{52} \cdot \frac{3}{51} \cdot \frac{2}{50} \cdot \frac{1}{49} \cdot \frac{48}{48} \cdot \binom{5}{1} = \frac{1}{54145}.$$

This is called the **conditional probability** because additional conditions are taken into account.

Conditional probability

- The **conditional probability** answers the question of how the probability of an event changes when we have additional information.

Conditional probability

- The **conditional probability** answers the question of how the probability of an event changes when we have additional information.
- The **conditional probability of A given B** is defined by

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad \text{with } P(B) > 0.$$

Conditional probability

- The **conditional probability** answers the question of how the probability of an event changes when we have additional information.
 - The **conditional probability of A given B** is defined by

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad \text{with } P(B) > 0.$$

- If B becomes the sample space we have $P(B|B) = 1$

Conditional probability

- The **conditional probability** answers the question of how the probability of an event changes when we have additional information.
 - The **conditional probability of A given B** is defined by

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad \text{with } P(B) > 0.$$

- If B becomes the sample space we have $P(B|B) = 1$
- Let A and B be **disjoint**, then $P(A \cap B) = 0$ and thus $P(A|B) = P(B|A) = 0$.

Conditional probability

- The **conditional probability** answers the question of how the probability of an event changes when we have additional information.
- The **conditional probability of A given B** is defined by

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \quad \text{with } P(B) > 0.$$

- If B becomes the sample space we have $P(B|B) = 1$
- Let A and B be **disjoint**, then $P(A \cap B) = 0$ and thus $P(A|B) = P(B|A) = 0$.

HW Draw two cards from a deck of cards. Let the events be D_1 = "first card is a diamond" and D_2 = "second card is a diamond". What are $P(D_2|D_1)$, $P(D_1|D_2)$ und $P(D_2|D_1^c)$?

Multiplication theorem and the Bayes theorem

- We rewrite the definition of conditional probability and obtain

$$P(A \cap B) = P(A|B) \cdot P(B) \quad \text{for } P(B) > 0 \quad \text{and}$$

$$P(A \cap B) = P(B|A) \cdot P(A) \quad \text{for } P(A) > 0.$$

Multiplication theorem and the Bayes theorem

- We rewrite the definition of conditional probability and obtain

$$P(A \cap B) = P(A|B) \cdot P(B) \quad \text{for } P(B) > 0 \quad \text{and}$$

$$P(A \cap B) = P(B|A) \cdot P(A) \quad \text{for } P(A) > 0.$$

... these equations are called **Multiplication theorem** ...

Multiplication theorem and the Bayes theorem

- We rewrite the definition of conditional probability and obtain

$$P(A \cap B) = P(A|B) \cdot P(B) \quad \text{for } P(B) > 0 \quad \text{and}$$

$$P(A \cap B) = P(B|A) \cdot P(A) \quad \text{for } P(A) > 0.$$

... these equations are called **Multiplication theorem** ...

- We equate the two expressions and obtain

$$P(A|B) = \frac{P(B|A) \cdot P(A)}{P(B)} \quad \text{and} \quad P(B|A) = \frac{P(A|B) \cdot P(B)}{P(A)}$$

which gives the formula for reversing ("turning around") conditional probabilities. The previous equation is called the **Bayes theorem** (for two events).

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are independent.

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are independent.

- Then,

$$\underline{P(A \cap B)} = P(A|B) \cdot P(B) = \underline{P(A) \cdot P(B)}$$

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are **independent**.

- Then,

$$\underline{P(A \cap B)} = P(A|B) \cdot P(B) = \underline{P(A) \cdot P(B)}$$

Thus, two events A and B are **independent** if it holds

$$P(A \cap B) = P(A) \cdot P(B)$$

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are **independent**.

- Then,

$$\underline{P(A \cap B)} = P(A|B) \cdot P(B) = \underline{P(A) \cdot P(B)}$$

Thus, two events A and B are **independent** if it holds

$$P(A \cap B) = P(A) \cdot P(B)$$

HW Let A and B be independent. Show that

- A and B^c
- A^c and B^c

are also independent.

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are **independent**.

- Then,

$$\underline{P(A \cap B)} = P(A|B) \cdot P(B) = \underline{P(A) \cdot P(B)}$$

Thus, two events A and B are **independent** if it holds

$$P(A \cap B) = P(A) \cdot P(B)$$

HW Let A and B be independent. Show that

- A and B^c ... Prove $P(A \cap B^c) = P(A) \cdot P(B^c)$
- A^c and B^c

are also independent.

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are **independent**.

- Then,

$$\underline{P(A \cap B)} = P(A|B) \cdot P(B) = \underline{P(A) \cdot P(B)}$$

Thus, two events A and B are **independent** if it holds

$$P(A \cap B) = P(A) \cdot P(B)$$

HW Let A and B be independent. Show that

- A and B^c ... Prove $P(A \cap B^c) = P(A) \cdot P(B^c)$
- A^c and B^c ... Prove $P(A^c \cap B^c) = P(A^c) \cdot P(B^c)$

are also independent.

Independence

- If for two events A und B it holds

$$P(A|B) = P(A)$$

then we say that A and B are **independent**.

- Then,

$$\underline{P(A \cap B)} = P(A|B) \cdot P(B) = \underline{P(A) \cdot P(B)}$$

Thus, two events A and B are **independent** if it holds

$$P(A \cap B) = P(A) \cdot P(B)$$

HW Let A and B be independent. Show that

- A and B^c ... Prove $P(A \cap B^c) = P(A) \cdot P(B^c)$
- A^c and B^c ... Prove $P(A^c \cap B^c) = P(A^c) \cdot P(B^c)$

are also independent.

HW Let A and B be independent. Knowing that $P(A|B) = 0.2$ and $P(B|A) = 0.5$, compute the probability $P(A^c \cap B)$.

Example

- We toss a fair coin four times. Consider the events
 $A = \text{"at least three heads"}$ and
 $B = \text{"first toss is tail"}$
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Example

- We toss a fair coin four times. Consider the events
 $A =$ "at least three heads" and
 $B =$ "first toss is tail"
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

Example

- We toss a fair coin four times. Consider the events
 $A = \text{"at least three heads"}$ and
 $B = \text{"first toss is tail"}$
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

- $\Omega = \{HHHH, HHHT, \dots, TTTT\}, \quad |\Omega| = 2^4 = 16$

Example

- We toss a fair coin four times. Consider the events
 $A = \text{"at least three heads"}$ and
 $B = \text{"first toss is tail"}$
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

- $\Omega = \{HHHH, HHHT, \dots, TTTT\}, \quad |\Omega| = 2^4 = 16$
- $A = \{HHHT, HHTH, HTHH, THHH, HHHH\}, \quad |A| = 5, \quad P(A) = \frac{5}{16}$

Example

- We toss a fair coin four times. Consider the events
 $A =$ "at least three heads" and
 $B =$ "first toss is tail"
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

- $\Omega = \{HHHH, HHHT, \dots, TTTT\}, \quad |\Omega| = 2^4 = 16$
- $A = \{HHHT, HHTH, HTHH, THHH, HHHH\}, \quad |A| = 5, \quad P(A) = \frac{5}{16}$
- $B = \{Tijk : i, j, k \in \{T, H\}\}, \quad |B| = 2^3 = 8, \quad P(B) = \frac{1}{2}$

Example

- We toss a fair coin four times. Consider the events
 $A =$ "at least three heads" and
 $B =$ "first toss is tail"
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

- $\Omega = \{HHHH, HHHT, \dots, TTTT\}, \quad |\Omega| = 2^4 = 16$
- $A = \{HHHT, HHTH, HTHH, THHH, HHHH\}, \quad |A| = 5, \quad P(A) = \frac{5}{16}$
- $B = \{Tijk : i, j, k \in \{T, H\}\}, \quad |B| = 2^3 = 8, \quad P(B) = \frac{1}{2}$
- $A \cap B = \{THHH\}, \quad |A \cap B| = 1, \quad P(A \cap B) = \frac{1}{16}$

Example

- We toss a fair coin four times. Consider the events
 $A = \text{"at least three heads"}$ and
 $B = \text{"first toss is tail"}$
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

- $\Omega = \{HHHH, HHHT, \dots, TTTT\}, \quad |\Omega| = 2^4 = 16$
- $A = \{HHHT, HHTH, HTHH, THHH, HHHH\}, \quad |A| = 5, \quad P(A) = \frac{5}{16}$
- $B = \{Tijk : i, j, k \in \{T, H\}\}, \quad |B| = 2^3 = 8, \quad P(B) = \frac{1}{2}$
- $A \cap B = \{THHH\}, \quad |A \cap B| = 1, \quad P(A \cap B) = \frac{1}{16}$
- The conditional probabilities are

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{16}}{\frac{1}{2}} = \frac{1}{8} \quad \text{and} \quad P(B|A) = \frac{P(B \cap A)}{P(A)} = \frac{\frac{1}{16}}{\frac{5}{16}} = \frac{1}{5}.$$

Example

- We toss a fair coin four times. Consider the events
 $A = \text{"at least three heads"}$ and
 $B = \text{"first toss is tail"}$
What are $P(A|B)$ and $P(B|A)$? Are A and B independent?

Answer:

- $\Omega = \{HHHH, HHHT, \dots, TTTT\}, \quad |\Omega| = 2^4 = 16$
- $A = \{HHHT, HHTH, HTHH, THHH, HHHH\}, \quad |A| = 5, \quad P(A) = \frac{5}{16}$
- $B = \{Tijk : i, j, k \in \{T, H\}\}, \quad |B| = 2^3 = 8, \quad P(B) = \frac{1}{2}$
- $A \cap B = \{THHH\}, \quad |A \cap B| = 1, \quad P(A \cap B) = \frac{1}{16}$
- The conditional probabilities are

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{16}}{\frac{1}{2}} = \frac{1}{8} \quad \text{and} \quad P(B|A) = \frac{P(B \cap A)}{P(A)} = \frac{\frac{1}{16}}{\frac{5}{16}} = \frac{1}{5}.$$

- Since $P(A|B) = \frac{1}{8} \neq \frac{5}{16} = P(A)$, we conclude that A and B are **not independent**.

Independence

- Three events A , B and C are **independent** if the following hold

Independence

- Three events A , B and C are **independent** if the following hold

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A \cap C) = P(A) \cdot P(C)$$

$$P(B \cap C) = P(B) \cdot P(C)$$

$$P(A \cap B \cap C) = P(A) \cdot P(B) \cdot P(C)$$

Independence

- Three events A , B and C are **independent** if the following hold

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A \cap C) = P(A) \cdot P(C)$$

$$P(B \cap C) = P(B) \cdot P(C)$$

$$P(A \cap B \cap C) = P(A) \cdot P(B) \cdot P(C)$$

HW Roll two dice and consider the following events

A = "first die is 3"

B = "the sum is 6"

C = "the sum is 7"

A is independent of :

- (a) B and C (b) only B (c) only C (d) neither B nor C

Independence

- Three events A , B and C are **independent** if the following hold

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A \cap C) = P(A) \cdot P(C)$$

$$P(B \cap C) = P(B) \cdot P(C)$$

$$P(A \cap B \cap C) = P(A) \cdot P(B) \cdot P(C)$$

HW Roll two dice and consider the following events

A = "first die is 3"

B = "the sum is 6"

C = "the sum is 7"

A is independent of :

- (a) B and C (b) only B (c) only C (d) neither B nor C

Answer: (c)

Law of total probability

- If only the conditional probabilities and the probabilities of the conditional event are known, the total probability of A results from

$$P(A) = P(A|B) \cdot P(B) + P(A|B^c) \cdot P(B^c),$$

where B^c denotes the opposite event to B .


Law of total probability

- If only the conditional probabilities and the probabilities of the conditional event are known, the total probability of A results from

$$P(A) = P(A|B) \cdot P(B) + P(A|B^c) \cdot P(B^c),$$

where B^c denotes the opposite event to B .

- Let C_1, C_2, \dots with $P(C_j) > 0$ be a **partition** of the sample space Ω , i.e. these events are pairwise disjoint $C_i \cap C_j = \emptyset$ for $i \neq j$ and $\bigcup_{i=1}^{\infty} C_i = \Omega$.




Law of total probability

- If only the conditional probabilities and the probabilities of the conditional event are known, the total probability of A results from

$$P(A) = P(A|B) \cdot P(B) + P(A|B^c) \cdot P(B^c),$$

where B^c denotes the opposite event to B .

- Let C_1, C_2, \dots with $P(C_j) > 0$ be a **partition** of the sample space Ω , i.e. these events are pairwise disjoint $C_i \cap C_j = \emptyset$ for $i \neq j$ and $\bigcup_{i=1}^{\infty} C_i = \Omega$.



- The **total probability** of an event A is given by

$$P(A) = \sum_{i=1}^{\infty} P(A \cap C_i) = \sum_{i=1}^{\infty} P(A|C_i) \cdot P(C_i)$$

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.
 - Let C_1, C_2, \dots be a partition of Ω and $P(C_i) > 0$ for all $i \geq 1$.
Then, for an event A with $P(A) > 0$ it holds:

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.
 - Let C_1, C_2, \dots be a partition of Ω and $P(C_i) > 0$ for all $i \geq 1$.
Then, for an event A with $P(A) > 0$ it holds:

$$P(C_i|A) = \frac{P(A \cap C_i)}{P(A)} = \frac{P(A|C_i) \cdot P(C_i)}{\sum_{j=1}^{\infty} P(A|C_j) \cdot P(C_j)}$$

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.
 - Let C_1, C_2, \dots be a partition of Ω and $P(C_i) > 0$ for all $i \geq 1$.
Then, for an event A with $P(A) > 0$ it holds:

$$P(C_i|A) = \frac{P(A \cap C_i)}{P(A)} = \frac{P(A|C_i) \cdot P(C_i)}{\sum_{j=1}^{\infty} P(A|C_j) \cdot P(C_j)}$$

The Bayes theorem
(for two and more events)

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.
 - Let C_1, C_2, \dots be a partition of Ω and $P(C_i) > 0$ for all $i \geq 1$.
Then, for an event A with $P(A) > 0$ it holds:

$$P(C_i|A) = \frac{P(A \cap C_i)}{P(A)} = \frac{P(A|C_i) \cdot P(C_i)}{\sum_{j=1}^{\infty} P(A|C_j) \cdot P(C_j)}$$

The Bayes theorem
(for two and more events)

- The partition events C_1, C_2, \dots are also called **hypotheses**

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.
 - Let C_1, C_2, \dots be a partition of Ω and $P(C_i) > 0$ for all $i \geq 1$.
Then, for an event A with $P(A) > 0$ it holds:

$$P(C_i|A) = \frac{P(A \cap C_i)}{P(A)} = \frac{P(A|C_i) \cdot P(C_i)}{\sum_{j=1}^{\infty} P(A|C_j) \cdot P(C_j)}$$

The Bayes theorem
(for two and more events)

- The partition events C_1, C_2, \dots are also called **hypotheses**
- $P(C_i)$ is called the **a priori** probability of C_i , i.e. the probability before the entry of A

The Bayes theorem

- The **Bayes theorem** describes the probability of an event based on prior knowledge of the conditions that might be related to the event.
 - Let C_1, C_2, \dots be a partition of Ω and $P(C_i) > 0$ for all $i \geq 1$.
Then, for an event A with $P(A) > 0$ it holds:

$$P(C_i|A) = \frac{P(A \cap C_i)}{P(A)} = \frac{P(A|C_i) \cdot P(C_i)}{\sum_{j=1}^{\infty} P(A|C_j) \cdot P(C_j)}$$

The Bayes theorem
(for two and more events)

- The partition events C_1, C_2, \dots are also called **hypotheses**
- $P(C_i)$ is called the **a priori** probability of C_i , i.e. the probability before the entry of A
- $P(C_i|A)$ is called the **posterior** probability, i.e. the probability after the entry of A

Example

- An urn contains 5 red and 2 green balls. A random ball is selected and replaced with a ball of the other color, then a second ball is drawn.
 - (1) What is the probability the second ball is red?
 - (2) What is the probability the first ball was red given the second ball was red?

Use the probability tree to solve the problems.

Answer:


- Denote by R_1 = "the first ball is red" and R_2 = "the second ball is red".
- R_1^c = "the first ball is green" and R_2^c = "the second ball is green"
- $P(R_1) = \frac{5}{7}$ and $P(R_1^c) = 1 - \frac{5}{7} = \frac{2}{7}$
- $P(R_2|R_1) = \frac{4}{7}$ and $P(R_2^c|R_1) = \frac{3}{7}$
- $P(R_2|R_1^c) = \frac{6}{7}$ and $P(R_2^c|R_1^c) = \frac{1}{7}$

Example cont.

- We draw the **probability tree**


Example cont.

- We draw the **probability tree**




Example cont.

- We draw the **probability tree**




Example cont.

- We draw the **probability tree**



Example cont.

- We draw the **probability tree**




- (1) From the Law of total probability we obtain

$$P(R_2) = P(R_2|R_1) \cdot P(R_1) + P(R_2|R_1^c) \cdot P(R_1^c) = \frac{5}{7} \cdot \frac{4}{7} + \frac{2}{7} \cdot \frac{6}{7} = \frac{32}{49}$$

Example cont.

- We draw the **probability tree**



- (1) From the Law of total probability we obtain

$$P(R_2) = P(R_2|R_1) \cdot P(R_1) + P(R_2|R_1^c) \cdot P(R_1^c) = \frac{5}{7} \cdot \frac{4}{7} + \frac{2}{7} \cdot \frac{6}{7} = \frac{32}{49}$$

- (2) By the Bayes theorem we obtain

$$P(R_1|R_2) = \frac{P(R_1 \cap R_2)}{P(R_2)} = \frac{P(R_2|R_1) \cdot P(R_1)}{P(R_2)} = \frac{20/49}{32/49} = \frac{20}{32}$$

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.
 - There are three doors, one door hides a car, two hide goats.

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.
 - There are three doors, one door hides a car, two hide goats.
 - The participant chooses any door.

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.
 - There are three doors, one door hides a car, two hide goats.
 - The participant chooses any door.
 - The presenter always opens a door with a goat (because he knows where the car is).

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.
 - There are three doors, one door hides a car, two hide goats.
 - The participant chooses any door.
 - The presenter always opens a door with a goat (because he knows where the car is).
 - The candidate can then change doors if he wants.

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.
 - There are three doors, one door hides a car, two hide goats.
 - The participant chooses any door.
 - The presenter always opens a door with a goat (because he knows where the car is).
 - The candidate can then change doors if he wants.

What is the best strategy for winning a car?

(a) switch (b) do not switch (c) it does not matter

Example: Monty Hall problem

- In a TW show, a candidate can win a car if he guesses correctly behind which of the three doors the car is.
 - There are three doors, one door hides a car, two hide goats.
 - The participant chooses any door.
 - The presenter always opens a door with a goat (because he knows where the car is).
 - The candidate can then change doors if he wants.

What is the best strategy for winning a car?

(a) switch (b) do not switch (c) it does not matter

Answer:

- We build a tree representing the switching strategy:
first the candidate chooses a door (then Monty shows a goat),
then the candidate switches doors or not

Example: Monty Hall problem


- Denote the doors by T_1 , T_2 and T_3 . Let the car be behind the first door T_1 .

Example: Monty Hall problem

- Denote the doors by T_1 , T_2 and T_3 . Let the car be behind the first door T_1 .
- We form the probability tree


Example: Monty Hall problem

- Denote the doors by T_1 , T_2 and T_3 . Let the car be behind the first door T_1 .
- We form the probability tree




Example: Monty Hall problem

- Denote the doors by T_1 , T_2 and T_3 . Let the car be behind the first door T_1 .
- We form the probability tree




Example: Monty Hall problem

- Denote the doors by T_1 , T_2 and T_3 . Let the car be behind the first door T_1 .
- We form the probability tree



Example: Monty Hall problem

- Denote the doors by T_1 , T_2 and T_3 . Let the car be behind the first door T_1 .
- We form the probability tree



- The probability of winning the car for both scenarios is

$$P(\text{win the car} | \text{stick to the first choice}) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}$$

$$P(\text{win the car} | \text{switch the first choice}) = \frac{1}{3} + \frac{1}{3} = \frac{2}{3}$$

Example: Batteries

- A container contains three types of batteries Type 1, Type 2 and Type 3 in the ratio of 20 : 30 : 50. A battery is good if it lasts more than 100 hours. The probability that a Type 1 battery is good is 0.7, the probability that a Type 2 battery is good is 0.4 and the probability that a Type 3 battery is good is 0.3. A battery is randomly taken from the container.
 - (1) What is the probability the chosen battery is good?
 - (2) What is the probability the battery is of Type 3 given it was not good?

Example: Batteries

- A container contains three types of batteries Type 1, Type 2 and Type 3 in the ratio of 20 : 30 : 50. A battery is good if it lasts more than 100 hours. The probability that a Type 1 battery is good is 0.7, the probability that a Type 2 battery is good is 0.4 and the probability that a Type 3 battery is good is 0.3. A battery is randomly taken from the container.
 - (1) What is the probability the chosen battery is good?
 - (2) What is the probability the battery is of Type 3 given it was not good?

Answer:

Example: Batteries

- A container contains three types of batteries Type 1, Type 2 and Type 3 in the ratio of 20 : 30 : 50. A battery is good if it lasts more than 100 hours. The probability that a Type 1 battery is good is 0.7, the probability that a Type 2 battery is good is 0.4 and the probability that a Type 3 battery is good is 0.3. A battery is randomly taken from the container.
 - (1) What is the probability the chosen battery is good?
 - (2) What is the probability the battery is of Type 3 given it was not good?

Answer:

- Let G = "battery is good", H_i = "battery is of Type i ", where $i = 1, 2, 3$.

Example: Batteries

- A container contains three types of batteries Type 1, Type 2 and Type 3 in the ratio of 20 : 30 : 50. A battery is good if it lasts more than 100 hours. The probability that a Type 1 battery is good is 0.7, the probability that a Type 2 battery is good is 0.4 and the probability that a Type 3 battery is good is 0.3. A battery is randomly taken from the container.
 - (1) What is the probability the chosen battery is good?
 - (2) What is the probability the battery is of Type 3 given it was not good?

Answer:

- Let G = "battery is good", H_i = "battery is of Type i ", where $i = 1, 2, 3$.

(1) Then,

$$\begin{aligned}P(G) &= P(G|H_1) \cdot P(H_1) + P(G|H_2) \cdot P(H_2) + P(G|H_3) \cdot P(H_3) \\&= 0.2 \cdot 0.7 + 0.3 \cdot 0.4 + 0.5 \cdot 0.3 = 0.41\end{aligned}$$

Example: Batteries

- A container contains three types of batteries Type 1, Type 2 and Type 3 in the ratio of 20 : 30 : 50. A battery is good if it lasts more than 100 hours. The probability that a Type 1 battery is good is 0.7, the probability that a Type 2 battery is good is 0.4 and the probability that a Type 3 battery is good is 0.3. A battery is randomly taken from the container.
 - What is the probability the chosen battery is good?
 - What is the probability the battery is of Type 3 given it was not good?

Answer:

- Let G = 'battery is good', H_i = "battery is of Type i ", where $i = 1, 2, 3$.

(1) Then,

$$\begin{aligned}P(G) &= P(G|H_1) \cdot P(H_1) + P(G|H_2) \cdot P(H_2) + P(G|H_3) \cdot P(H_3) \\&= 0.2 \cdot 0.7 + 0.3 \cdot 0.4 + 0.5 \cdot 0.3 = 0.41\end{aligned}$$

(2)

$$P(H_3|G^c) = \frac{P(G^c|H_3) \cdot P(H_3)}{P(G^c)} = \frac{P(G^c|H_3) \cdot P(H_3)}{1 - P(G)} = \frac{0.5 \cdot 0.7}{0.59} = \frac{35}{59} = 0.593$$

More examples

HW

A blood test reacts 95% positive if there is a disease. It also reacts to 1% "false positive". It is assumed that 0.5% of the population are ill.

- (1) What is the probability a randomly selected person whose blood test is positive is actually sick?
- (2) What is the probability of being sick if the blood test is positive?

HW

The Randomizer holds the 6-sided die in one fist and the 8-sided die in the other. The Roller selects one of the Randomizer's fists and takes the die face down. The Roller rolls the die in secret and reports the result to the table.

- Given the reported number, what is the probability that the 6-sided die was selected? (Find the probability for each possible number reported.)

Questions

A few multiple-choice questions

- (1) A medical treatment has a success rate of 0.8. Two patients will be treated with this treatment. Assuming the results are independent for the two patients, what is the probability that at least one of them will be successfully cured?
- (a) 0.96
 - (b) 0.32
 - (c) 0.64
 - (d) 0.04
- (2) In a list of 15 households, 9 own homes and 6 do not own homes. Four households are randomly selected from these 15 households. Find the probability that the number of households in these four who own homes is at most one.
- (a) 0.1536
 - (b) 0.1792
 - (c) 0.3456
 - (d) 0.4752

A few multiple-choice questions

- (3) Suppose that there are 4 women and 8 men. How many 5 person committees can be formed with exactly 2 women and 3 men?

- (a) $\binom{12}{5}$
- (b) $\binom{4}{2} \cdot \binom{8}{2}$
- (c) $\binom{4}{2} \cdot \binom{8}{3}$
- (d) $\binom{12}{5} - \binom{4}{1} \cdot \binom{8}{4}$

- (4) Suppose box A contains 4 red and 5 blue coins and box B contains 6 red and 3 blue coins. A coin is chosen at random from the box A and placed in box B. Finally, a coin is chosen at random from among those now in box B. What is the probability a blue coin was transferred from box A to box B given that the coin chosen from box B is red?

- (a) $\frac{15}{29}$
- (b) $\frac{14}{29}$
- (c) $\frac{1}{2}$
- (d) $\frac{7}{10}$

A few multiple-choice questions

- (5) A fair die is rolled. Find the probability of getting an even number or a number bigger than 2.
- (a) $\frac{1}{3}$
 - (b) $\frac{7}{12}$
 - (c) $\frac{2}{3}$
 - (d) $\frac{5}{6}$
- (6) Anna has an unfair coin. She knows that when flipping this coin, head is obtained twice as often as a tail. Anna rolls this coin until she obtains a tail. What is the probability that she rolled the coin at most three times?
- (a) $\frac{19}{27}$
 - (b) $\frac{26}{27}$
 - (c) $\frac{5}{9}$
 - (d) $\frac{4}{27}$

Thank you for your attention!