


Surrounding the one-sample t -test



All examples are fictitious. All data are simulated and the graphics were created with the statistical program package R.

The materials are protected by copyright and are only provided for personal use for studies at TU Vienna. Further use is not permitted. In particular, it is not permitted to distribute the materials or make them publicly available (e.g. in social networks, on learning platforms, etc.).

Sämtliche Beispiele sind frei erfunden. Alle Daten sind simuliert und die Grafiken wurden mit statistischen Programmpaket R erstellt.


Die Materialien sind urheberrechtlich geschützt und dürfen ausschließlich für den Eigengebrauch im Rahmen des Studiums an der TU Wien genutzt werden. Eine weitere Nutzung ist nicht gestattet. Insbesondere ist es nicht gestattet, die Materialien zu verbreiten oder öffentlich zugänglich zu machen (etwa im Rahmen sozialer Netzwerke, Lernplattformen etc.).

Reminder

- How is the **mean** distributed under normal distribution?


Reminder

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$




Reminder

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$




Reminder

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed




Reminder

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)



Reminder


- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}



Reminder

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)

Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$



From the z -test to the t -test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)

Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters

From the z-test to the t-test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)

Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters
- Concept in z-Test:
 - μ fixed via null hypothesis: $H_0 : \mu = \mu_0$ holds true

From the z -test to the t -test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)

Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters
- Concept in z -Test:
 - μ fixed via null hypothesis: $H_0 : \mu = \mu_0$ holds true
 - σ assumed as known. But unknown in practice (Problem!)

From the z-test to the t-test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters
- Concept in z-Test:
 - μ fixed via null hypothesis: $H_0 : \mu = \mu_0$ holds true
 - σ assumed as known. But unknown in practice (Problem!)Way out: *Estimate* σ through the empirical standard deviation S

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

From the z-test to the t-test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters
- Concept in z-Test:
 - μ fixed via null hypothesis: $H_0 : \mu = \mu_0$ holds true
 - σ assumed as known. But unknown in practice (Problem!)Way out: *Estimate* σ through the empirical standard deviation S

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- Plug in estimator S

$$\frac{\bar{X} - \mu}{S / \sqrt{n}}$$

From the z-test to the t-test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters
- Concept in z-Test:
 - μ fixed via null hypothesis: $H_0 : \mu = \mu_0$ holds true
 - σ assumed as known. But unknown in practice (Problem!)Way out: *Estimate* σ through the empirical standard deviation S

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- Plug in estimator S

$$\frac{\bar{X} - \mu}{S / \sqrt{n}} \sim t(n-1)$$

t-distributed with $n-1$ degrees of freedom.

From the z -test to the t -test

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- Point of view in statistics: μ and σ are unknown population parameters
- Concept in z -Test:
 - μ fixed via null hypothesis: $H_0 : \mu = \mu_0$ holds true
 - σ assumed as known. But unknown in practice (Problem!)Way out: *Estimate* σ through the empirical standard deviation S

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

- Plug in estimator S

$$\frac{\bar{X} - \mu}{S / \sqrt{n}} \sim t(n-1)$$

t-distributed with $n - 1$ degrees of freedom. What is the *t*-distribution?

t -distribution


- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S / \sqrt{n}} \sim t(n - 1)$$

- $N(0, 1)$



t -distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$


$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

- $N(0,1)$

- $t(1)$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$

Γ denotes the Gamma function

t-distribution


- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

- $N(0,1)$
- $t(1)$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n ,


Γ denotes the Gamma function

t -distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0,1) \quad \text{vs} \quad \frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

- $N(0,1)$
- $t(2)$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n ,

Γ denotes the Gamma function

t -distribution


- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

- $N(0,1)$
- $t(10)$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n ,

Γ denotes the Gamma function

t -distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$


$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$

• $N(0,1)$

• $t(20)$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n ,

Γ denotes the Gamma function


t -distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n ,

Γ denotes the Gamma function


t-distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n ,
- $t(n)$ symmetric around 0

Γ denotes the Gamma function


t-distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n , Γ denotes the Gamma function
- $t(n)$ symmetric around 0
- $t(n)$ has heavier tails (polynomial) than the $N(0, 1)$ -distribution (exponential)


t -distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n , Γ denotes the Gamma function
- $t(n)$ symmetric around 0
- $t(n)$ has heavier tails (polynomial) than the $N(0, 1)$ -distribution (exponential)
 - Intuition: estimation of σ via S increases the variability of the rescaled mean


t-distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}, x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n , Γ denotes the Gamma function
- $t(n)$ symmetric around 0
- $t(n)$ has heavier tails (polynomial) than the $N(0, 1)$ -distribution (exponential)
 - Intuition: estimation of σ via S increases the variability of the rescaled mean
- the density of $t(n)$ converges pointwise to the density of $N(0, 1)$


t-distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) :\Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}$, $x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n , Γ denotes the Gamma function
- $t(n)$ symmetric around 0
- $t(n)$ has heavier tails (polynomial) than the $N(0, 1)$ -distribution (exponential)
 - Intuition: estimation of σ via S increases the variability of the rescaled mean
- the density of $t(n)$ converges pointwise to the density of $N(0, 1)$
 - Intuition: the estimation of σ via S gets more precise (law of large numbers)


t-distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S/\sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) \Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}$, $x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n , Γ denotes the Gamma function
- $t(n)$ symmetric around 0
- $t(n)$ has heavier tails (polynomial) than the $N(0, 1)$ -distribution (exponential)
 - Intuition: estimation of σ via S increases the variability of the rescaled mean
- the density of $t(n)$ converges pointwise to the density of $N(0, 1)$
 - Intuition: the estimation of σ via S gets more precise (law of large numbers)
- in R via `rt(..., df=n)`, `dt(..., df=n)`, etc. with `df` denoting 'degrees of freedom'


t-distribution

- Let X_1, \dots, X_n be i.i.d. random variables and $X_1 \sim N(\mu, \sigma^2)$

$$\frac{\bar{X} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$$

vs

$$\frac{\bar{X} - \mu}{S / \sqrt{n}} \sim t(n-1)$$



The $t(n)$ -distribution ($n \in \{1, 2, \dots\}$)

- Definition: $X \sim t(n) \Leftrightarrow X$ has density $f(x) = \frac{\Gamma((n+1)/2)}{\sqrt{n\pi} \cdot \Gamma(n/2)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}$, $x \in \mathbb{R}$
- $t(n)$ has a one parameter: degrees of freedom n , Γ denotes the Gamma function
- $t(n)$ symmetric around 0
- $t(n)$ has heavier tails (polynomial) than the $N(0, 1)$ -distribution (exponential)
 - Intuition: estimation of σ via S increases the variability of the rescaled mean
- the density of $t(n)$ converges pointwise to the density of $N(0, 1)$
 - Intuition: the estimation of σ via S gets more precise (law of large numbers)
- in R via `rt(..., df=n)`, `dt(..., df=n)`, etc. with `df` denoting 'degrees of freedom'


From what is said, it is (hopefully) plausible, but not rigorously proven, that $\frac{\bar{X} - \mu}{S / \sqrt{n}} \sim t(n-1)$.

For details see e.g., Messer, M. and Schneider, G. *Statistik: Theorie und Praxis im Dialog*, Springer Berlin

Example from last lecture

Reminder:


- Runtimes of an algorithm implemented by $n = 16$ students that took a certain programming class.



Example from last lecture

Reminder:

- Runtimes of an algorithm implemented by $n = 16$ students that took a certain programming class.



- Delicate assertion from a colleague of the lecturer: " The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been $\mu_0 = 30$

From z -test to the t -test

- Assertion about a huge 'population'

From z-test to the t -test

- Assertion about a huge 'population'
- the null hypothesis, that the mean runtime was $\mu_0 = 30$, was judged by the z-test (and it was rejected)

From z-test to the *t*-test

- Assertion about a huge 'population'
- the null hypothesis, that the mean runtime was $\mu_0 = 30$, was judged by the z-test (and it was rejected)
- now same procedure, but replace the z-statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$

(σ assumed as known)

From z-test to the *t*-test

- Assertion about a huge 'population'
- the null hypothesis, that the mean runtime was $\mu_0 = 30$, was judged by the z-test (and it was rejected)
- now same procedure, but replace the z-statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$

(σ assumed as known)
through the *t*-statistic

$$T := \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

(unknown σ estimated via $S \rightarrow$ practically applicable)

From z-test to the t -test

- Assertion about a huge 'population'
- the null hypothesis, that the mean runtime was $\mu_0 = 30$, was judged by the z-test (and it was rejected)
- now same procedure, but replace the z-statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$

(σ assumed as known)
through the t -statistic

$$T := \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

(unknown σ estimated via $S \rightarrow$ practically applicable)

We then speak about the (one-sample) t -test

From z-test to the *t*-test

- Assertion about a huge 'population'
- the null hypothesis, that the mean runtime was $\mu_0 = 30$, was judged by the z-test (and it was rejected)
- now same procedure, but replace the z-statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma / \sqrt{n}}$$

(σ assumed as known)
through the *t*-statistic

$$T := \frac{\bar{X} - \mu_0}{S / \sqrt{n}}$$

(unknown σ estimated via $S \rightarrow$ practically applicable)

We then speak about the (one-sample) *t*-test

- Note that T is a proper *statistic* in the sense that it is a bare function of the random variables / the data. Particularly, it does not depend on unknown parameters

The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)


The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)

The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)
- *Test statistic* for the evaluation of the data (measures discrepancy). Now t -statistic

$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$



The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)


- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)

- *Test statistic* for the evaluation of the data (measures discrepancy). Now t -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if H_0 holds true*

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$



The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)

- *Test statistic* for the evaluation of the data (measures discrepancy). Now t -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$


- *Distribution of the test statistic if H_0 holds true*

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *p -value:* quantifies discrepancy (judge t according to the distribution of T)

$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if H_0 holds true (here: two-sided Test)



The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)

- *Test statistic* for the evaluation of the data (measures discrepancy). Now t -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if H_0 holds true*


$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *p-value:* quantifies discrepancy (judge t according to the distribution of T)

$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if H_0 holds true (here: two-sided Test)

- *Decision:* Reject the null hypothesis, because $p \leq \alpha$
Say: the observed discrepancy was *significant* ($p < 10^{-4}$)



The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)

- *Test statistic* for the evaluation of the data (measures discrepancy). Now t -statistic

$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if H_0 holds true*


$$T = \frac{\bar{X} - \mu_0}{S / \sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *p -value:* quantifies discrepancy (judge t according to the distribution of T)

$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if H_0 holds true (here: two-sided Test)

- *Decision:* Reject the null hypothesis, because $p \leq \alpha \Leftrightarrow t \in R$
Say: the observed discrepancy was *significant* ($p < 10^{-4}$)



The (one-sample) t -Test according to 'Student' (google: W.S. Gosset)

- *Set significance level:* Choose (e.g.,) $\alpha = 5\%$
- *Model assumption:* X_1, \dots, X_n i.i.d. RVs, with $X_1 \sim N(\mu, \sigma^2)$, $\mu \in \mathbb{R}$, $\sigma > 0$ ($n = 16$)
(The data x_1, \dots, x_n are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation μ and unknown variance σ^2)
- *Null hypothesis:* $H_0 : \mu = 30$
Describes the assertion: the claimed expectation is $\mu_0 = 30$)

- *Test statistic* for the evaluation of the data (measures discrepancy). Now t -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if H_0 holds true*

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *p -value:* quantifies discrepancy (judge t according to the distribution of T)


$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if H_0 holds true (here: two-sided Test)

- *Decision:* Reject the null hypothesis, because $p \leq \alpha \Leftrightarrow t \in R$


Say: the observed discrepancy was *significant* ($p < 10^{-4}$)

- *Interpretation:* If H_0 holds true, then something very unlikely was observed. In that sense, the data are hardly compatible with H_0 .




t -test vs. z -test

- 'Result', interpretation etc. are in the t -test the 'same' as in the z -test. Differences are only in the test statistics and in the distributions




$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$




$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

t -test vs. z -test

- 'Result', interpretation etc. are in the t -test the 'same' as in the z -test. Differences are only in the the test statistics and in the distributions
- We take a closer look:




$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$




$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

t-test vs. *z*-test

- 'Result', interpretation etc. are in the *t*-test the 'same' as in the *z*-test. Differences are only in the test statistics and in the distributions
- We take a closer look:
 - $s = 8.7$ (*t*-test) has underestimated $\sigma = 11$ (*z*-test)




$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$




$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

t -test vs. z -test

- 'Result', interpretation etc. are in the t -test the 'same' as in the z -test. Differences are only in the test statistics and in the distributions
- We take a closer look:
 - $s = 8.7$ (t -test) has underestimated $\sigma = 11$ (z -test)
 - Thus, $t \approx -6.2$ more extreme than $z \approx -4.9$ (as s and σ in the denominator)




$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$




$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

t -test vs. z -test

- 'Result', interpretation etc. are in the t -test the 'same' as in the z -test. Differences are only in the test statistics and in the distributions
- We take a closer look:
 - $s = 8.7$ (t -test) has underestimated $\sigma = 11$ (z -test)
 - Thus, $t \approx -6.2$ more extreme than $z \approx -4.9$ (as s and σ in the denominator)
 - However: p -value in t -test ($\approx 1.5 \cdot 10^{-5}$) larger than p in z -test ($\approx 9 \cdot 10^{-7}$)




$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$




$$z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

t-test vs. *z*-test

- 'Result', interpretation etc. are in the *t*-test the 'same' as in the *z*-test. Differences are only in the test statistics and in the distributions
- We take a closer look:
 - $s = 8.7$ (*t*-test) has underestimated $\sigma = 11$ (*z*-test)
 - Thus, $t \approx -6.2$ more extreme than $z \approx -4.9$ (as s and σ in the denominator)
 - However: p -value in *t*-test ($\approx 1.5 \cdot 10^{-5}$) larger than p in *z*-test ($\approx 9 \cdot 10^{-7}$)
 - No contradiction, because the tails of the *t*-distribution are heavier than those of $N(0, 1)$ (visually not observable in the graphics)




$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$




$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11/4} \approx -4.9$$

t -test vs. z -test

- 'Result', interpretation etc. are in the t -test the 'same' as in the z -test. Differences are only in the test statistics and in the distributions
- We take a closer look:
 - $s = 8.7$ (t -test) has underestimated $\sigma = 11$ (z -test)
 - Thus, $t \approx -6.2$ more extreme than $z \approx -4.9$ (as s and σ in the denominator)
 - However: p -value in t -test ($\approx 1.5 \cdot 10^{-5}$) larger than p in z -test ($\approx 9 \cdot 10^{-7}$)
 - No contradiction, because the tails of the t -distribution are heavier than those of $N(0, 1)$ (visually not observable in the graphics)
 - For the same reason: in the t -test slightly smaller rejection area R : for $\alpha = 5\%$ it holds first regarding $t(15)$ that $q_{\alpha/2} = -2.13$, and second regarding $N(0, 1)$ that $q_{\alpha/2} = -1.96$ (while $q_{\alpha/2}$ denotes the $\alpha/2$ -quantiles)



$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} \approx \frac{16.5 - 30}{8.7 / 4} \approx -6.2$$



$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \approx \frac{16.5 - 30}{11 / 4} \approx -4.9$$

The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \mathbb{V}ar(X_1) \in (0, \infty)$

The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \mathbb{V}ar(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$

The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \mathbb{V}ar(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

(n=16)



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$


(n=16)



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean


$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean


$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean


$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$




The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

- In the context of statistics: σ unknown population parameter



The standard error of the mean


- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

- In the context of statistics: σ unknown population parameter
- Estimate σ via S

<u>Definition:</u> $SEM := \frac{S}{\sqrt{n}}$
--

Standard Error of the Mean



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean


$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

- In the context of statistics: σ unknown population parameter
- Estimate σ via S

<u>Definition:</u> $SEM := \frac{S}{\sqrt{n}}$
--

Standard Error of the Mean

- Meaning: the estimated variability of the mean



The standard error of the mean

- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$
- Mean $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- Standard deviation of the mean

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

- In the context of statistics: σ unknown population parameter
- Estimate σ via S

<u>Definition:</u> $SEM := \frac{S}{\sqrt{n}}$
--

Standard Error of the Mean

- Meaning: the estimated variability of the mean
- The t -statistic measures the discrepancy $\bar{X} - \mu_0$ in the units SEM

$T = \frac{\bar{X} - \mu_0}{SEM}$

The standard error of the mean


- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$

$$SEM = \frac{S}{\sqrt{n}}$$

- The t -statistic measures the discrepancy $\bar{X} - \mu_0$ in the units SEM

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

Under H_0 : $T \sim t(n-1)$



- Intuition: If $H_0 : \mu = \mu_0$ holds true

- $|t| = 1 \Leftrightarrow |\bar{x} - \mu_0| = 1 \cdot sem$

it is not unlikely to observe a discrepancy of one sem

The standard error of the mean


- Let X_1, \dots, X_n be i.i.d. RVs and $\sigma^2 = \text{Var}(X_1) \in (0, \infty)$

$$SEM = \frac{S}{\sqrt{n}}$$


- The t -statistic measures the discrepancy $\bar{X} - \mu_0$ in the units SEM

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

Under $H_0: T \sim t(n-1)$



Under $H_0: T \sim t(n-1)$



- Intuition: If $H_0: \mu = \mu_0$ holds true

- $|t| = 1 \Leftrightarrow |\bar{x} - \mu_0| = 1 \cdot \text{sem}$

it is not unlikely to observe a discrepancy of one sem

- $|t| = 5 \Leftrightarrow |\bar{x} - \mu_0| = 5 \cdot \text{sem}$

it is very unlikely to observe a discrepancy of five sem


'chance has a hard time realizing this'

From the standard error to the confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

$$SEM = \frac{S}{\sqrt{n}}$$



- In our example: $t = \frac{\bar{x} - \mu_0}{sem} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$


it is very unlikely to observe a discrepancy $|\bar{x} - \mu_0|$ of $6.2 \cdot sem$ under H_0 .

From the standard error to the confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$ with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

$$SEM = \frac{S}{\sqrt{n}}$$




$$(*) \quad \alpha = \mathbb{P}_{H_0}(T \in R)$$

From the standard error to the confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$ with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

$$SEM = \frac{S}{\sqrt{n}}$$



$$\begin{aligned} (*) \quad \alpha &= \mathbb{P}_{H_0}(T \in R) \\ &= \mathbb{P}_{H_0}(|\bar{X} - \mu_0| \geq q_{1-\alpha/2} \cdot SEM) \end{aligned}$$

($q_{1-\alpha/2}$ is the $(1 - \alpha/2)$ -quantile of $t(n - 1)$)



From the standard error to the confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$ with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

$$SEM = \frac{S}{\sqrt{n}}$$

runtimes



(*)

$$\alpha = \mathbb{P}_{H_0}(T \in R)$$

$$= \mathbb{P}_{H_0}(|\bar{X} - \mu_0| \geq q_{1-\alpha/2} \cdot SEM)$$

$$= \mathbb{P}_{H_0}((\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM) \not\ni \mu_0)$$

($q_{1-\alpha/2}$ is the $(1 - \alpha/2)$ -quantile of $t(n - 1)$)



From the standard error to the confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$ with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$

$$T = \frac{\bar{X} - \mu_0}{SEM}$$

$$SEM = \frac{S}{\sqrt{n}}$$

runtimes



$$\begin{aligned}
 (*) \quad \alpha &= \mathbb{P}_{H_0}(T \in R) \\
 &= \mathbb{P}_{H_0}(|\bar{X} - \mu_0| \geq q_{1-\alpha/2} \cdot SEM) \\
 &= \mathbb{P}_{H_0}((\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM) \not\ni \mu_0) \\
 &\quad (q_{1-\alpha/2} \text{ is the } (1 - \alpha/2)\text{-quantile of } t(n - 1))
 \end{aligned}$$


- Meaning: Under H_0 , the interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$

From the standard error to the confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$ with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$



$$\begin{aligned}
 (*) \quad \alpha &= \mathbb{P}_{H_0}(T \in R) \\
 &= \mathbb{P}_{H_0}(|\bar{X} - \mu_0| \geq q_{1-\alpha/2} \cdot SEM) \\
 &= \mathbb{P}_{H_0}((\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM) \not\supset \mu_0) \\
 &\quad (q_{1-\alpha/2} \text{ is the } (1 - \alpha/2)\text{-quantile of } t(n - 1))
 \end{aligned}$$

- Meaning: Under H_0 , the interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$

- I is called a $(1 - \alpha)$ -confidence interval for μ (abbreviate: C.I.)

Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n - 1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$

Confidence interval


- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval


$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$

Under $H_0: T \sim t(n-1)$



Under $H_0: T \sim t(n-1)$





Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- $t \in R$
Reject H_0



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- $t \in R \Leftrightarrow i \not\in \mu_0$

(see $(*)$ in previous slide)

Reject H_0 if and only if μ_0 is not overlapped by the confidence interval i



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- $t \in R \Leftrightarrow i \not\in \mu_0$

(see (*) in previous slide)

Reject H_0 if and only if μ_0 is not overlapped by the confidence interval i

- Meaning: Equivalence of test and confidence interval (\rightarrow 'Student's C.I.')



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- Decrease of α
 \leftrightarrow increase of $q_{1-\alpha/2}$



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- Decrease of α
 - \Leftrightarrow increase of $q_{1-\alpha/2}$
 - \Leftrightarrow increase of the width of the confidence interval



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- Decrease of α
 - \Leftrightarrow increase of $q_{1-\alpha/2}$
 - \Leftrightarrow increase of the width of the confidence interval



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- Decrease of α
 - \leftrightarrow increase of $q_{1-\alpha/2}$
 - \leftrightarrow increase of the width of the confidence interval



Confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs and $X_1 \sim N(\mu, \sigma^2)$, with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $t(n-1)$

Under $H_0 : \mu = \mu_0$, the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability $1 - \alpha$



- Decrease of α


\leftrightarrow increase of $q_{1-\alpha/2}$

\leftrightarrow increase of the width of the confidence interval

Plausible: 'If μ_0 shall be overlapped with large probability, then I must be large'

Interpretation


- Delicate assertion from a colleague of the lecturer: " The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been $\mu_0 = 30'$



- The 95% confidence interval (here $\alpha = 5\%$) does not overlap μ_0

Interpretation


- Delicate assertion from a colleague of the lecturer: " The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been $\mu_0 = 30'$



- The 95% confidence interval (here $\alpha = 5\%$) does not overlap μ_0
- If the colleague is right, then the data describe an unlikely observation. In fact, one of the those observations that appear only in 5% of cases, if he is right.

Interpretation


- Delicate assertion from a colleague of the lecturer: " The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been $\mu_0 = 30'$



- The 95% confidence interval (here $\alpha = 5\%$) does not overlap μ_0
- If the colleague is right, then the data describe an unlikely observation. In fact, one of the those observations that appear only in 5% of cases, if he is right.
- In that sense the data are barely compatible with the assertion


Interpretation

- Delicate assertion from a colleague of the lecturer: " The course was held by the lecturer a couple of times before. If all participants that have ever taken the course had implemented this algorithm, then the mean runtime would have been $\mu_0 = 30'$




- The 95% confidence interval (here $\alpha = 5\%$) does not overlap μ_0
- If the colleague is right, then the data describe an unlikely observation. In fact, one of the those observations that appear only in 5% of cases, if he is right.
- In that sense the data are barely compatible with the assertion
- Intuitively: $q_{1-\alpha/2} \approx 2.13 \rightarrow$ the c.i. has a diameter of about $4sem$ \rightarrow the distance $|\bar{X} - \mu_0|$ is about $6sem$. That is a large distance if we take in account that the typical deviation of \bar{X} from μ_0 is about $1SEM$

Data not bell-shaped




- Here: $n = 36$ data

Data not bell-shaped




- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$

Data not bell-shaped




- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$
- but distribution of the data is asymmetric

Data not bell-shaped




- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$
- but distribution of the data is asymmetric
- Particularly, not approximately bell-shaped

Data not bell-shaped



- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$
- but distribution of the data is asymmetric
- Particularly, not approximately bell-shaped
- Previous model assumption, that the data are sampled from the normal distribution is not appropriate


Data not bell-shaped



- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$
- but distribution of the data is asymmetric
- Particularly, not approximately bell-shaped
- Previous model assumption, that the data are sampled from the normal distribution is not appropriate
- Way out: asymptotic normality of the mean

Reminder


- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)Interpretation: the typical deviation of the **mean** from its expectation is σ/\sqrt{n}



Reminder

- How is the **mean** distributed under normal distribution?
- Let X_1, \dots, X_n be independent and identically distributed (i.i.d.) random variables and $X_1 \sim N(\mu, \sigma^2)$
- For the mean it holds $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$
 - \bar{X} is also normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$ (equal to the expectation of X_i)
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma / \sqrt{n}$ (decrease of factor $1/\sqrt{n}$)

Interpretation: the typical deviation of the **mean** from its expectation is σ / \sqrt{n}
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?


Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \stackrel{d}{\rightarrow} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)



Asymptotic normality of the mean


- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)



Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)


(n=36)



Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)


(n=36)



Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)


(n=36)



Asymptotic normality of the mean


- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \stackrel{d}{\rightarrow} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)

(n=36)




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \stackrel{d}{\rightarrow} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \stackrel{d}{\rightarrow} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \stackrel{d}{\rightarrow} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)




Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)



Asymptotic normality of the mean

- How is the **mean** distributed for a large sample size n ?
- Let X_1, \dots, X_n i.i.d. random variables and $\text{Var}(X_1) \in (0, \infty)$ (not necessarily normally distributed!)
- Then $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{d}{\approx} N(\mu, \sigma^2/n)$ approximately
 - \bar{X} is approximately normally distributed
 - \bar{X} has expectation $\mu_{\bar{X}} = \mu$
 - \bar{X} has standard deviation $\sigma_{\bar{X}} = \sigma/\sqrt{n}$
- **Standardization:** $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \xrightarrow{d} N(0, 1)$ for $n \rightarrow \infty$ (central limit theorem)



Asymptotic one-sample test and confidence interval

- Let X_1, \dots, X_n be i.i.d. RVs with $\mu \in \mathbb{R}$ and $\text{Var}(X_1) \in (0, \infty)$, and let $q_{1-\alpha/2}$ be the $(1 - \alpha/2)$ -quantile of $N(0, 1)$

Under $H_0 : \mu = \mu_0$ it approximatively holds for large n

$$T = \frac{\bar{X} - \mu_0}{SEM} \stackrel{d}{\approx} N(0, 1)$$


and equivalently: the confidence interval

$$I := (\bar{X} - q_{1-\alpha/2} \cdot SEM, \bar{X} + q_{1-\alpha/2} \cdot SEM)$$

overlaps the parameter μ_0 with probability about $1 - \alpha$


- i.e.: test and confidence interval are constructed according to 'Student'. Only difference: Use the quantiles of $N(0, 1)$, instead of $t(n - 1)$

Approximate procedure




- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$

Approximate procedure



- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$
- Approximate 95%-confidence interval does not overlap μ_0


Approximate procedure



- Here: $n = 36$ data
- $H_0 : \mu = \mu_0$
- Approximate 95%-confidence interval does not overlap μ_0
→ reject H_0 on the 5% level

Question


Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?



Here t -test naively:

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?




Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?




Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)
- $s \approx 3$ (bell-shaped distribution, 2/3 of the data captured)

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?




Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)
- $s \approx 3$ (bell-shaped distribution, 2/3 of the data captured)
- $sem \approx 3/10$ (100 data points)

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?




Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)
- $s \approx 3$ (bell-shaped distribution, 2/3 of the data captured)
- $sem \approx 3/10$ (100 data points)
- $q \approx 2$ (n large, 97.5%-quantile of $N(0, 1)$ is ≈ 1.96)

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?




Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)
- $s \approx 3$ (bell-shaped distribution, 2/3 of the data captured)
- $sem \approx 3/10$ (100 data points)
- $q \approx 2$ (n large, 97.5%-quantile of $N(0, 1)$ is ≈ 1.96)
- Together: C.I. $i \approx (4.4, 5.6)$ ($\bar{x} \pm q \cdot sem$)

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?




Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)
- $s \approx 3$ (bell-shaped distribution, 2/3 of the data captured)
- $sem \approx 3/10$ (100 data points)
- $q \approx 2$ (n large, 97.5%-quantile of $N(0, 1)$ is ≈ 1.96)
- Together: C.I. $i \approx (4.4, 5.6)$ ($\bar{x} \pm q \cdot sem$)
- Hence: Reject H_0 (C.I. does not overlap μ_0)

Question

Can $H_0 : \mu = \mu_0 = 3$ be rejected on the 5%-level?



Here t -test naively:

- $\bar{x} \approx 5$ (balance in equilibrium)
- $s \approx 3$ (bell-shaped distribution, 2/3 of the data captured)
- $sem \approx 3/10$ (100 data points)
- $q \approx 2$ (n large, 97.5%-quantile of $N(0, 1)$ is ≈ 1.96)
- Together: C.I. $i \approx (4.4, 5.6)$ ($\bar{x} \pm q \cdot sem$)
- Hence: Reject H_0 (C.I. does not overlap μ_0)

Of course, this is not 'precise', but the important message is that \bar{x} is further than $6sem$ away from μ_0 . This is extremely far! More precise estimates would have yielded the same message!

t-test in R

```
# Enter data
x <- c(...)
# perform t-test
t.test(x,mu=3,...)
# Output
```

One Sample t-test

```
data:  x
t = 8.3512, df = 99, p-value = 4.22e-13
alternative hypothesis: true mean is not equal to 3
95 percent confidence interval:
 4.745578 5.833572
sample estimates:
mean of x
5.289575
```

- Our naive estimates were plausible
 - Mean and C.I. fit well
 - \bar{x} is even further than $8sem$ apart from μ_0

Thank you!