







# Revision



All examples are fictitious. All data are simulated and the graphics were created with the statistical program package R.


The materials are protected by copyright and are only provided for personal use for studies at TU Vienna. Further use is not permitted. In particular, it is not permitted to distribute the materials or make them publicly available (e.g. in social networks, on learning platforms, etc.).

Sämtliche Beispiele sind frei erfunden. Alle Daten sind simuliert und die Grafiken wurden mit statistischen Programmpaket R erstellt.

Die Materialien sind urheberrechtlich geschützt und dürfen ausschließlich für den Eigengebrauch im Rahmen des Studiums an der TU Wien genutzt werden. Eine weitere Nutzung ist nicht gestattet. Insbesondere ist es nicht gestattet, die Materialien zu verbreiten oder öffentlich zugänglich zu machen (etwa im Rahmen sozialer Netzwerke, Lernplattformen etc.).

# Overview

We differentiate:  
Probability theory  
(Stochastics)  
=  
Theory of randomness



# Overview

We differentiate:

Probability theory  
(Stochastics)

=



Theory of randomness

and

Statistics

=

Description of data  $\longrightarrow$   
(using stochastic **models**)



# Data collection

How long is the runtime of an algorithm that you implemented?

# Data collection

How long is the runtime of an algorithm that you implemented?

$n = 121$  students requested (same technical setup)

# Data collection

How long is the runtime of an algorithm that you implemented?

$n = 121$  students requested (same technical setup)

Results (in seconds):

24.6, 24, 31.4, 29.9, 37.8, 19.9, 46.1, 32.8, 30.3, 29, 47.1, 27.8, 33.8, 30.1, 53.3, 23.8, 32.1, 4.2, 42.8, 25.2, 52.3, 35, 30.1, 43.2, 25.4, 62.5, 35.4, 25.2, 37.6, 37.1, 22.9, 29.5, 44.5, 34.8, 33.3, 21.9, 37.2, 24, 37, 34, 24.1, 10.8, 24.9, 37.2, 52, 30.8, 22, 18.6, 22, 26.8, 52.3, 27, 23.6, 33.5, 30.8, 20.9, 35.6, 37.2, 57.5, 46.2, 36.1, 19.8, 38.1, 36.9, 26.5, 23.6, 30.3, 49.9, 39, 50.2, 35.7, 11.4, 24.1, 27.5, 36.4, 29.8, 49, 42.6, 22.5, 32.7, 34.3, 21.4, 34.7, 47.3, 20.3, 35.4, 41.8, 24.9, 15.2, 42.2, 29.1, 25.1, 22.7, 41, 28.2, 30.3, 25.6, 41.8, 16.6, 38, 43.1, 29.5, 40.3, 20.5, 39.9, 24.5, 33.7, 14.6, 23.3, 36.7, 34.7, 34.9, 39.1, 32.2, 43, 12.1, 19.8, 27.4, 39.3, 35, 46.3

# Data collection

How long is the runtime of an algorithm that you implemented?

$n = 121$  students requested (same technical setup)

Results (in seconds):

24.6, 24, 31.4, 29.9, 37.8, 19.9, 46.1, 32.8, 30.3, 29, 47.1, 27.8, 33.8, 30.1, 53.3, 23.8, 32.1, 4.2, 42.8, 25.2, 52.3, 35, 30.1, 43.2, 25.4, 62.5, 35.4, 25.2, 37.6, 37.1, 22.9, 29.5, 44.5, 34.8, 33.3, 21.9, 37.2, 24, 37, 34, 24.1, 10.8, 24.9, 37.2, 52, 30.8, 22, 18.6, 22, 26.8, 52.3, 27, 23.6, 33.5, 30.8, 20.9, 35.6, 37.2, 57.5, 46.2, 36.1, 19.8, 38.1, 36.9, 26.5, 23.6, 30.3, 49.9, 39, 50.2, 35.7, 11.4, 24.1, 27.5, 36.4, 29.8, 49, 42.6, 22.5, 32.7, 34.3, 21.4, 34.7, 47.3, 20.3, 35.4, 41.8, 24.9, 15.2, 42.2, 29.1, 25.1, 22.7, 41, 28.2, 30.3, 25.6, 41.8, 16.6, 38, 43.1, 29.5, 40.3, 20.5, 39.9, 24.5, 33.7, 14.6, 23.3, 36.7, 34.7, 34.9, 39.1, 32.2, 43, 12.1, 19.8, 27.4, 39.3, 35, 46.3

We see:  $n$  data:  $x_1 = 24.6, x_2 = 24.0, \dots, x_n = 46.3$



# Data collection

How long is the runtime of an algorithm that you implemented?

$n = 121$  students requested (same technical setup)

Results (in seconds):

24.6, 24, 31.4, 29.9, 37.8, 19.9, 46.1, 32.8, 30.3, 29, 47.1, 27.8, 33.8, 30.1, 53.3, 23.8, 32.1, 4.2, 42.8, 25.2, 52.3, 35, 30.1, 43.2, 25.4, 62.5, 35.4, 25.2, 37.6, 37.1, 22.9, 29.5, 44.5, 34.8, 33.3, 21.9, 37.2, 24, 37, 34, 24.1, 10.8, 24.9, 37.2, 52, 30.8, 22, 18.6, 22, 26.8, 52.3, 27, 23.6, 33.5, 30.8, 20.9, 35.6, 37.2, 57.5, 46.2, 36.1, 19.8, 38.1, 36.9, 26.5, 23.6, 30.3, 49.9, 39, 50.2, 35.7, 11.4, 24.1, 27.5, 36.4, 29.8, 49, 42.6, 22.5, 32.7, 34.3, 21.4, 34.7, 47.3, 20.3, 35.4, 41.8, 24.9, 15.2, 42.2, 29.1, 25.1, 22.7, 41, 28.2, 30.3, 25.6, 41.8, 16.6, 38, 43.1, 29.5, 40.3, 20.5, 39.9, 24.5, 33.7, 14.6, 23.3, 36.7, 34.7, 34.9, 39.1, 32.2, 43, 12.1, 19.8, 27.4, 39.3, 35, 46.3

We see:  $n$  data:  $x_1 = 24.6, x_2 = 24.0, \dots, x_n = 46.3$

We understand: nothing?

# Data collection

How long is the runtime of an algorithm that you implemented?

$n = 121$  students requested (same technical setup)

Results (in seconds):

24.6, 24, 31.4, 29.9, 37.8, 19.9, 46.1, 32.8, 30.3, 29, 47.1, 27.8, 33.8, 30.1, 53.3, 23.8, 32.1, 4.2, 42.8, 25.2, 52.3, 35, 30.1, 43.2, 25.4, 62.5, 35.4, 25.2, 37.6, 37.1, 22.9, 29.5, 44.5, 34.8, 33.3, 21.9, 37.2, 24, 37, 34, 24.1, 10.8, 24.9, 37.2, 52, 30.8, 22, 18.6, 22, 26.8, 52.3, 27, 23.6, 33.5, 30.8, 20.9, 35.6, 37.2, 57.5, 46.2, 36.1, 19.8, 38.1, 36.9, 26.5, 23.6, 30.3, 49.9, 39, 50.2, 35.7, 11.4, 24.1, 27.5, 36.4, 29.8, 49, 42.6, 22.5, 32.7, 34.3, 21.4, 34.7, 47.3, 20.3, 35.4, 41.8, 24.9, 15.2, 42.2, 29.1, 25.1, 22.7, 41, 28.2, 30.3, 25.6, 41.8, 16.6, 38, 43.1, 29.5, 40.3, 20.5, 39.9, 24.5, 33.7, 14.6, 23.3, 36.7, 34.7, 34.9, 39.1, 32.2, 43, 12.1, 19.8, 27.4, 39.3, 35, 46.3

We see:  $n$  data:  $x_1 = 24.6, x_2 = 24.0, \dots, x_n = 46.3$

We understand: nothing?



Thus: descriptive Statistics  $\rightarrow$  graphical representation and summary of data

# Descriptive statistics

- Graphical representations
- Summarizing statistics



# Graphical representations

- stripchart





# Graphical representations

- stripchart
- histogram





# Graphical representations

- stripchart
- histogram
- boxplot





# Graphical representations

- stripchart
- histogram
- boxplot
- barplot



# Graphical representations

- stripchart
- histogram
- boxplot
- barplot
- plot






# Graphical representations

- stripchart
- histogram
- boxplot
- barplot
- plot
- ...?


# Summary statistics

- mean  $\bar{x}$




# Summary statistics

- mean  $\bar{x}$
- empirical standard deviation  $s$  (and emp. variance  $s^2$ )




# Summary statistics

- mean  $\bar{x}$
- empirical standard deviation  $s$  (and emp. variance  $s^2$ )
- minimum  $x_{(1)}$  and maximum  $x_{(n)}$



# Summary statistics

- mean  $\bar{x}$
- empirical standard deviation  $s$  (and emp. variance  $s^2$ )
- minimum  $x_{(1)}$  and maximum  $x_{(n)}$
- empirical quantiles  $q_p$  (e.g., **median**, **quartiles**)



# Summary statistics


- mean  $\bar{x}$
- empirical standard deviation  $s$  (and emp. variance  $s^2$ )
- minimum  $x_{(1)}$  and maximum  $x_{(n)}$
- empirical quantiles  $q_p$  (e.g., **median**, **quartiles**)
- ...?

For details see the lecture: 'Descriptive Statistics'


# Statistical hypothesis test

- How compatible are the data with an assertion?

runtimes




Null hypothesis rejected




# Statistical hypothesis test

- How **compatible** are the **data** with an **assertion**?
- Notion of **incompatibility** via probability statements within statistical models ('data interpreted as realizations of random variables')

**runtimes**



Null hypothesis rejected







# Statistical hypothesis test

- How **compatible** are the **data** with an **assertion**?
- Notion of **incompatibility** via probability statements within statistical models ('data interpreted as realizations of random variables')
- Formalized in hypothesis tests (or confidence intervals)

runtimes




Null hypothesis rejected




# Statistical hypothesis test

- How **compatible** are the **data** with an **assertion**?
- Notion of **incompatibility** via probability statements within statistical models ('data interpreted as realizations of random variables')
- Formalized in hypothesis tests (or confidence intervals)
- Need additional statistics
  - **test statistics** ( $t$ -statistic,  $f$ -statistic,  $\chi^2$ -statistic etc.)
  - **auxiliary statistics** (e.g., standard errors)
  - **$p$ -value**

runtimes




Null hypothesis rejected




# Statistical hypothesis test

- How **compatible** are the **data** with an **assertion**?
- Notion of **incompatibility** via probability statements within statistical models ('data interpreted as realizations of random variables')
- Formalized in hypothesis tests (or confidence intervals)
- Need additional statistics
  - **test statistics** ( $t$ -statistic,  $f$ -statistic,  $\chi^2$ -statistic etc.)
  - **auxiliary statistics** (e.g., standard errors)
  - **$p$ -value**

runtimes



Null hypothesis rejected



For general concepts of modeling and testing see the lecture: 'Basic ideas of hypothesis testing'

## Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )


## Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

# Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )
- *Test statistic* for the evaluation of the data (measures discrepancy). Now  $t$ -statistic

$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$



# Example: the (one-sample) $t$ -Test


- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

- *Test statistic* for the evaluation of the data (measures discrepancy). Now  $t$ -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if  $H_0$  holds true*

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$



# Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

- *Test statistic* for the evaluation of the data (measures discrepancy). Now  $t$ -statistic

$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$


- *Distribution of the test statistic if  $H_0$  holds true*

$$T = \frac{\bar{X} - \mu_0}{S / \sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *$p$ -value:* quantifies discrepancy (judge  $t$  according to the distribution of  $T$ )

$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Under  $H_0$ :  $T \sim t(15)$



Probability to make an observation which is at least as extreme as in the data, if  $H_0$  holds true (here: two-sided test)



# Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

- *Test statistic* for the evaluation of the data (measures discrepancy). Now  $t$ -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if  $H_0$  holds true*


$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *$p$ -value:* quantifies discrepancy (judge  $t$  according to the distribution of  $T$ )

$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if  $H_0$  holds true (here: two-sided test)

- *Decision:* Reject the null hypothesis, because  $p \leq \alpha$   
Say: the observed discrepancy was *significant* ( $p < 10^{-4}$ )



# Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

- *Test statistic* for the evaluation of the data (measures discrepancy). Now  $t$ -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- Distribution of the test statistic if  $H_0$  holds true


$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- $p$ -value: quantifies discrepancy (judge  $t$  according to the distribution of  $T$ )

$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if  $H_0$  holds true (here: two-sided test)

- *Decision:* Reject the null hypothesis, because  $p \leq \alpha \Leftrightarrow t \in R$   
Say: the observed discrepancy was *significant* ( $p < 10^{-4}$ )



# Example: the (one-sample) $t$ -Test

- *Set significance level:* Choose (e.g.,)  $\alpha = 5\%$
- *Model assumption:*  $X_1, \dots, X_n$  i.i.d. RVs, with  $X_1 \sim N(\mu, \sigma^2)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$  ( $n = 16$ )  
(The data  $x_1, \dots, x_n$  are assumed to be realizations of i.i.d. normal distributed RVs with unknown expectation  $\mu$  and unknown variance  $\sigma^2$ )
- *Null hypothesis:*  $H_0 : \mu = 30$   
Describes the assertion: the claimed expectation is  $\mu_0 = 30$ )

- *Test statistic* for the evaluation of the data (measures discrepancy). Now  $t$ -statistic

$$t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}} \approx \frac{16.5 - 30}{8.7/4} \approx -6.2$$

- *Distribution of the test statistic if  $H_0$  holds true*

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \stackrel{H_0}{\sim} t(15)$$

- *$p$ -value:* quantifies discrepancy (judge  $t$  according to the distribution of  $T$ )


$$p = \mathbb{P}_{H_0}(|T| \geq |t|) \approx 1.5 \cdot 10^{-5}$$

Probability to make an observation which is at least as extreme as in the data, if  $H_0$  holds true (here: two-sided test)



- *Decision:* Reject the null hypothesis, because  $p \leq \alpha \Leftrightarrow t \in R$

Say: the observed discrepancy was *significant* ( $p < 10^{-4}$ )

- *Interpretation:* If  $H_0$  holds true, then something very unlikely was observed. In that sense, the data are hardly compatible with  $H_0$ .



# Equivalence of test and confidence interval





Reject  $H_0$  if and only if  $\mu_0$  is not overlapped by the confidence interval

$$i := (\bar{x} - q_{1-\alpha/2} \cdot \text{sem}, \bar{x} + q_{1-\alpha/2} \cdot \text{sem})$$

with  $\text{sem} := s / \sqrt{n}$

# Equivalence of test and confidence interval



Reject  $H_0$  if and only if  $\mu_0$  is not overlapped by the confidence interval  
 $i := (\bar{x} - q_{1-\alpha/2} \cdot \text{sem}, \bar{x} + q_{1-\alpha/2} \cdot \text{sem})$  with  $\text{sem} := s / \sqrt{n}$



Recall that the upper test and confidence interval are referred to as 'Student's' versions (one-sample situation).

But also remember that the structure of the test statistic and the confidence interval is inherited to other situations, i.e., 'more general'

$$T = \frac{\spadesuit - \clubsuit}{\heartsuit} \quad \text{and} \quad I = (\spadesuit - q \cdot \heartsuit, \spadesuit + q \cdot \heartsuit)$$

while  $\spadesuit$  denotes some summary statistic,  $\clubsuit$  denotes the null-parameter, and  $\heartsuit$  a standard-error of the statistic, see e.g., the two-sample situation.

# Equivalence of test and confidence interval



Reject  $H_0$  if and only if  $\mu_0$  is not overlapped by the confidence interval  
 $i := (\bar{x} - q_{1-\alpha/2} \cdot \text{sem}, \bar{x} + q_{1-\alpha/2} \cdot \text{sem})$  with  $\text{sem} := s / \sqrt{n}$


Recall that the upper test and confidence interval are referred to as 'Student's' versions (one-sample situation).

But also remember that the structure of the test statistic and the confidence interval is inherited to other situations, i.e., 'more general'


$$T = \frac{\spadesuit - \clubsuit}{\heartsuit} \quad \text{and} \quad I = (\spadesuit - q \cdot \heartsuit, \spadesuit + q \cdot \heartsuit)$$

while  $\spadesuit$  denotes some summary statistic,  $\clubsuit$  denotes the null-parameter, and  $\heartsuit$  a standard-error of the statistic, see e.g., the two-sample situation.  
 For details see the lecture: 'Surrounding the one-sample t-test'

# Two-sample $t$ -test




# Two-sample $t$ -test



- Let  $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$  be independent RVs with  
 $X_i \sim N(\mu_1, \sigma_1^2)$  for  $i = 1, \dots, n_1$  and  
 $Y_j \sim N(\mu_2, \sigma_2^2)$  for  $j = 1, \dots, n_2$  with  $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
- $d = \mu_2 - \mu_1$
- Let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $t(\nu)$ -distribution (R knows  $\nu$ )



# Two-sample $t$ -test




- Let  $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$  be independent RVs with  
 $X_i \sim N(\mu_1, \sigma_1^2)$  for  $i = 1, \dots, n_1$  and  
 $Y_j \sim N(\mu_2, \sigma_2^2)$  for  $j = 1, \dots, n_2$  with  $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
- $d = \mu_2 - \mu_1$
- Let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $t(\nu)$ -distribution (R knows  $\nu$ )

Under  $H_0 : d = d_0$  it holds (approx)

$$T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$$

# Two-sample $t$ -test



- Let  $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$  be independent RVs with  
 $X_i \sim N(\mu_1, \sigma_1^2)$  for  $i = 1, \dots, n_1$  and  
 $Y_j \sim N(\mu_2, \sigma_2^2)$  for  $j = 1, \dots, n_2$  with  $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
- $d = \mu_2 - \mu_1$
- Let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $t(\nu)$ -distribution (R knows  $\nu$ )

Under  $H_0 : d = d_0$  it holds (approx)



$$T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$$

and equivalently: The confidence interval

$$I := \left( (\bar{Y} - \bar{X}) - q_{1-\alpha/2} \cdot \sqrt{SEM_y^2 + SEM_x^2}, (\bar{Y} - \bar{X}) + q_{1-\alpha/2} \cdot \sqrt{SEM_y^2 + SEM_x^2} \right)$$

overlaps the parameter  $d_0$  with probability (approx)  $1 - \alpha$

# Two-sample $t$ -test



- Let  $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$  be independent RVs with  
 $X_i \sim N(\mu_1, \sigma_1^2)$  for  $i = 1, \dots, n_1$  and  
 $Y_j \sim N(\mu_2, \sigma_2^2)$  for  $j = 1, \dots, n_2$  with  $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
- $d = \mu_2 - \mu_1$
- Let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $t(\nu)$ -distribution (R knows  $\nu$ )

Under  $H_0 : d = d_0$  it holds (approx)

$$T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$$



and equivalently: The confidence interval

$$I := \left( (\bar{Y} - \bar{X}) - q_{1-\alpha/2} \cdot \sqrt{SEM_y^2 + SEM_x^2}, (\bar{Y} - \bar{X}) + q_{1-\alpha/2} \cdot \sqrt{SEM_y^2 + SEM_x^2} \right)$$

overlaps the parameter  $d_0$  with probability (approx)  $1 - \alpha$

- Under  $H_0 : d = 5$  ( $\Leftrightarrow \mu_2 = \mu_1 + 5$ )  $\rightarrow$  here: can not reject  $H_0$  ( $\rightarrow$  Welch-test)

# Two-sample $t$ -test



- Let  $X_1, \dots, X_{n_1}, Y_1, \dots, Y_{n_2}$  be independent RVs with  
 $X_i \sim N(\mu_1, \sigma_1^2)$  for  $i = 1, \dots, n_1$  and  
 $Y_j \sim N(\mu_2, \sigma_2^2)$  for  $j = 1, \dots, n_2$  with  $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2) \in \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} \times \mathbb{R}^+$
- $d = \mu_2 - \mu_1$
- Let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $t(\nu)$ -distribution (R knows  $\nu$ )

Under  $H_0 : d = d_0$  it holds (approx)

$$T := \frac{(\bar{Y} - \bar{X}) - d_0}{\sqrt{SEM_y^2 + SEM_x^2}} \sim t(\nu)$$

and equivalently: The confidence interval

$$I := \left( (\bar{Y} - \bar{X}) - q_{1-\alpha/2} \cdot \sqrt{SEM_y^2 + SEM_x^2}, (\bar{Y} - \bar{X}) + q_{1-\alpha/2} \cdot \sqrt{SEM_y^2 + SEM_x^2} \right)$$

overlaps the parameter  $d_0$  with probability (approx)  $1 - \alpha$


- Under  $H_0 : d = 5$  ( $\Leftrightarrow \mu_2 = \mu_1 + 5$ )  $\rightarrow$  here: can not reject  $H_0$  ( $\rightarrow$  Welch-test)

For details see the lecture: 'Surrounding the two-sample  $t$ -test'

# ANOVA

Model:



- Let  $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$  be independent RVs and for  $i = 1, \dots, k$  let  $X_{i,j} \sim N(\mu_i, \sigma^2)$  for  $j = 1, \dots, n_i$ , with  $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus,  $k$  groups and  $n = \sum_{i=1}^k n_i$  observations in total



# ANOVA

## Model:

- Let  $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$  be independent RVs and for  $i = 1, \dots, k$  let  $X_{i,j} \sim N(\mu_i, \sigma^2)$  for  $j = 1, \dots, n_i$ , with  $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus,  $k$  groups and  $n = \sum_{i=1}^k n_i$  observations in total




# ANOVA

Model:

- Let  $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$  be independent RVs and for  $i = 1, \dots, k$  let  $X_{i,j} \sim N(\mu_i, \sigma^2)$  for  $j = 1, \dots, n_i$ , with  $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus,  $k$  groups and  $n = \sum_{i=1}^k n_i$  observations in total

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} = \frac{\text{'variability between the groups'}}{\text{'variability within the groups'}}$$




# ANOVA

Model:


- Let  $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$  be independent RVs and for  $i = 1, \dots, k$  let  $X_{i,j} \sim N(\mu_i, \sigma^2)$  for  $j = 1, \dots, n_i$ , with  $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus,  $k$  groups and  $n = \sum_{i=1}^k n_i$  observations in total

Under  $H_0 : \mu_1 = \dots = \mu_k$  it holds

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i,\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i,\cdot})^2} \sim \mathcal{F}(k-1, n-k)$$



- Here: reject  $H_0$ .






# ANOVA

Model:

- Let  $X_{1,1}, \dots, X_{1,n_1}, X_{2,1}, \dots, X_{2,n_2}, \dots, X_{k,1}, \dots, X_{k,n_k}$  be independent RVs and for  $i = 1, \dots, k$  let  $X_{i,j} \sim N(\mu_i, \sigma^2)$  for  $j = 1, \dots, n_i$ , with  $(\mu_1, \dots, \mu_k, \sigma^2) \in \mathbb{R}^k \times \mathbb{R}^+$
- Thus,  $k$  groups and  $n = \sum_{i=1}^k n_i$  observations in total


Under  $H_0 : \mu_1 = \dots = \mu_k$  it holds

$$F := \frac{\frac{1}{k-1} \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X})^2}{\frac{1}{n-k} \sum_{i,j} (X_{i,j} - \bar{X}_{i\cdot})^2} \sim \mathcal{F}(k-1, n-k)$$



- Here: reject  $H_0$ .

Under  $H_0: F \sim \mathcal{F}(k-1, n-k)$



For details on comparisons on multiple groups see the lecture: 'Analysis of variance and multiple testing'


# Linear regression

- For  $i = 1, \dots, n$  let

$$Y_i = \beta_0 + \beta_1 \cdot x_i + \sigma Z_i,$$

with  $Z_1, \dots, Z_n$  i.i.d. RVs and  $Z_1 \sim N(0, 1)$ , and  $(\beta_0, \beta_1, \sigma) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^+$

- $q_{1-\alpha/2}$  the  $(1 - \alpha/2)$ -quantile of the  $t(n - 2)$ -distribution



# Linear regression


- For  $i = 1, \dots, n$  let

$$Y_i = \beta_0 + \beta_1 \cdot x_i + \sigma Z_i,$$

with  $Z_1, \dots, Z_n$  i.i.d. RVs and  $Z_1 \sim N(0, 1)$ , and  $(\beta_0, \beta_1, \sigma) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^+$

- $q_{1-\alpha/2}$  the  $(1 - \alpha/2)$ -quantile of the  $t(n - 2)$ -distribution
- estimators  $B_0, B_1$  and  $S_r$  via 'least-squares',  $SE_{B_1} = S_r / (s_x \cdot \sqrt{n - 1})$

**n = 20**



# Linear regression


- For  $i = 1, \dots, n$  let

$$Y_i = \beta_0 + \beta_1 \cdot x_i + \sigma Z_i,$$


with  $Z_1, \dots, Z_n$  i.i.d. RVs and  $Z_1 \sim N(0, 1)$ , and  $(\beta_0, \beta_1, \sigma) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^+$

- $q_{1-\alpha/2}$  the  $(1 - \alpha/2)$ -quantile of the  $t(n - 2)$ -distribution
- estimators  $B_0, B_1$  and  $S_r$  via 'least-squares',  $SE_{B_1} = S_r / (s_x \cdot \sqrt{n - 1})$

**n = 20**



$H_0: \beta_1 = 0$



# Linear regression

- For  $i = 1, \dots, n$  let

$$Y_i = \beta_0 + \beta_1 \cdot x_i + \sigma Z_i,$$

with  $Z_1, \dots, Z_n$  i.i.d. RVs and  $Z_1 \sim N(0, 1)$ , and  $(\beta_0, \beta_1, \sigma) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^+$

- $q_{1-\alpha/2}$  the  $(1 - \alpha/2)$ -quantile of the  $t(n - 2)$ -distribution
- estimators  $B_0, B_1$  and  $S_r$  via 'least-squares',  $SE_{B_1} = S_r / (s_x \cdot \sqrt{n - 1})$

Under  $H_0 : \beta_1 = \beta_1^{(0)}$  it holds

$$T := \frac{B_1 - \beta_1^{(0)}}{SE_{B_1}} \sim t(n - 2)$$

and equivalently: the confidence interval

$$I := (B_1 - q_{1-\alpha/2} \cdot SE_{B_1}, B_1 + q_{1-\alpha/2} \cdot SE_{B_1})$$

overlaps  $\beta_1^{(0)}$  with probability  $1 - \alpha$

# Linear regression



- For  $i = 1, \dots, n$  let

$$Y_i = \beta_0 + \beta_1 \cdot x_i + \sigma Z_i,$$

with  $Z_1, \dots, Z_n$  i.i.d. RVs and  $Z_1 \sim N(0, 1)$ , and  $(\beta_0, \beta_1, \sigma) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^+$

- $q_{1-\alpha/2}$  the  $(1 - \alpha/2)$ -quantile of the  $t(n - 2)$ -distribution
- estimators  $B_0, B_1$  and  $S_r$  via 'least-squares',  $SE_{B_1} = S_r / (s_x \cdot \sqrt{n - 1})$

**n = 20**



- Here: reject  $H_0 : \beta_1 = 0$

# Linear regression



- For  $i = 1, \dots, n$  let

$$Y_i = \beta_0 + \beta_1 \cdot x_i + \sigma Z_i,$$

with  $Z_1, \dots, Z_n$  i.i.d. RVs and  $Z_1 \sim N(0, 1)$ , and  $(\beta_0, \beta_1, \sigma) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^+$

- $q_{1-\alpha/2}$  the  $(1 - \alpha/2)$ -quantile of the  $t(n - 2)$ -distribution
- estimators  $B_0, B_1$  and  $S_r$  via 'least-squares',  $SE_{B_1} = S_r / (s_x \cdot \sqrt{n - 1})$

**n = 20**




- Here: reject  $H_0 : \beta_1 = 0$

For details on correlation and linear regression see the lecture: 'Linear Regression'

# Frequencies

- Let  $Y_{1,1}, \dots, Y_{1,n_1}, Y_{2,1}, \dots, Y_{2,n_2}$  be independent RVs with  $Y_{1,i} \sim \text{ber}(p_1)$  for  $i = 1, \dots, n_1$  and  $Y_{2,j} \sim \text{ber}(p_2)$  for  $j = 1, \dots, n_2$ ,  
and  $(p_1, p_2) \in (0, 1)^2$
- let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $N(0, 1)$ -distribution



$$H_j := \frac{1}{n_j} \sum_{i=1}^{n_j} Y_{j,i} \quad \text{and} \quad SE_{H_j} := \sqrt{\frac{H_j(1 - H_j)}{n_j}}$$



# Frequencies

- Let  $Y_{1,1}, \dots, Y_{1,n_1}, Y_{2,1}, \dots, Y_{2,n_2}$  be independent RVs with  $Y_{1,i} \sim \text{ber}(p_1)$  for  $i = 1, \dots, n_1$  and  $Y_{2,j} \sim \text{ber}(p_2)$  for  $j = 1, \dots, n_2$ ,  
and  $(p_1, p_2) \in (0, 1)^2$
- let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $N(0, 1)$ -distribution

Under  $H_0 : p_2 - p_1 = 0$  it holds (approximately)

$$Z := \frac{(\textcolor{red}{H}_2 - \textcolor{red}{H}_1) - 0}{\sqrt{\textcolor{brown}{SE}_{H_2}^2 + \textcolor{brown}{SE}_{H_1}^2}} \stackrel{d}{\approx} N(0, 1)$$

and equivalently: the confidence interval



$$I := \left( (\textcolor{red}{H}_2 - \textcolor{red}{H}_1) - q_{1-\alpha/2} \cdot \sqrt{\textcolor{brown}{SE}_{H_2}^2 + \textcolor{brown}{SE}_{H_1}^2}, (\textcolor{red}{H}_2 - \textcolor{red}{H}_1) + q_{1-\alpha/2} \cdot \sqrt{\textcolor{brown}{SE}_{H_2}^2 + \textcolor{brown}{SE}_{H_1}^2} \right)$$

overlaps 0 with probability about  $1 - \alpha$

$$\textcolor{red}{H}_j := \frac{1}{n_j} \sum_{i=1}^{n_j} Y_{j,i} \quad \text{and} \quad \textcolor{brown}{SE}_{H_j} := \sqrt{\frac{H_j(1 - H_j)}{n_j}}$$

# Frequencies

- Let  $Y_{1,1}, \dots, Y_{1,n_1}, Y_{2,1}, \dots, Y_{2,n_2}$  be independent RVs with  $Y_{1,i} \sim \text{ber}(p_1)$  for  $i = 1, \dots, n_1$  and  $Y_{2,j} \sim \text{ber}(p_2)$  for  $j = 1, \dots, n_2$ ,  
and  $(p_1, p_2) \in (0, 1)^2$
- let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $N(0, 1)$ -distribution





$$H_j := \frac{1}{n_j} \sum_{i=1}^{n_j} Y_{j,i} \quad \text{and}$$

$$SE_{H_j} := \sqrt{\frac{H_j(1 - H_j)}{n_j}}$$

# Frequencies

- Let  $Y_{1,1}, \dots, Y_{1,n_1}, Y_{2,1}, \dots, Y_{2,n_2}$  be independent RVs with  $Y_{1,i} \sim \text{ber}(p_1)$  for  $i = 1, \dots, n_1$  and  $Y_{2,j} \sim \text{ber}(p_2)$  for  $j = 1, \dots, n_2$ ,  
and  $(p_1, p_2) \in (0, 1)^2$
- let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $N(0, 1)$ -distribution





$$H_j := \frac{1}{n_j} \sum_{i=1}^{n_j} Y_{j,i} \quad \text{and}$$

$$SE_{H_j} := \sqrt{\frac{H_j(1 - H_j)}{n_j}}$$

- Here: can not reject  $H_0 : p_2 = p_1$

# Frequencies

- Let  $Y_{1,1}, \dots, Y_{1,n_1}, Y_{2,1}, \dots, Y_{2,n_2}$  be independent RVs with  $Y_{1,i} \sim \text{ber}(p_1)$  for  $i = 1, \dots, n_1$  and  $Y_{2,j} \sim \text{ber}(p_2)$  for  $j = 1, \dots, n_2$ ,  
and  $(p_1, p_2) \in (0, 1)^2$
- let  $q_{1-\alpha/2}$  be the  $(1 - \alpha/2)$ -quantile of the  $N(0, 1)$ -distribution



$$H_j := \frac{1}{n_j} \sum_{i=1}^{n_j} Y_{j,i} \quad \text{and} \quad SE_{H_j} := \sqrt{\frac{H_j(1 - H_j)}{n_j}}$$


- Here: can not reject  $H_0 : p_2 = p_1$

For details on frequencies see the lecture: 'Proportions'

# The $\chi^2$ -test (goodness of fit)

- Let  $\mathfrak{X} = (X_1, \dots, X_d)^t \sim \text{mult}(n, p)$ , with  $p \in (0, 1)^d$  and  $\sum_{k=1}^d p_k = 1$

observed frequencies



# The $\chi^2$ -test (goodness of fit)


- Let  $\mathfrak{X} = (\mathbf{X}_1, \dots, \mathbf{X}_d)^t \sim \text{mult}(n, p)$ , with  $p \in (0, 1)^d$  and  $\sum_{k=1}^d p_k = 1$

Under  $H_0 : p = (p_{0,1}, \dots, p_{0,d})^t$  it holds (approximately)

$$X^2 := \sum_{k=1}^d \frac{(\mathbf{X}_k - \mathbb{E}_{H_0}[\mathbf{X}_k])^2}{\mathbb{E}_{H_0}[\mathbf{X}_k]} \stackrel{d}{\approx} \chi^2(d-1)$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2(d-1)$  as  $n \rightarrow \infty$

- here:  $n = 120$  and  $d = 6$ , as well as  $p_0 = (1/6, \dots, 1/6)^t$



# The $\chi^2$ -test (goodness of fit)



- Let  $\mathfrak{X} = (X_1, \dots, X_d)^t \sim \text{mult}(n, p)$ , with  $p \in (0, 1)^d$  and  $\sum_{k=1}^d p_k = 1$

Under  $H_0 : p = (p_{0,1}, \dots, p_{0,d})^t$  it holds (approximately)

$$X^2 := \sum_{k=1}^d \frac{(X_k - \mathbb{E}_{H_0}[X_k])^2}{\mathbb{E}_{H_0}[X_k]} \stackrel{d}{\approx} \chi^2(d-1)$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2(d-1)$  as  $n \rightarrow \infty$

- here:  $n = 120$  and  $d = 6$ , as well as  $p_0 = (1/6, \dots, 1/6)^t$
- Let  $q_{1-\alpha}$  denote the  $(1 - \alpha)$ -quantile of the  $\chi^2(d-1)$ -distribution



# The $\chi^2$ -test (goodness of fit)



- Let  $\mathfrak{X} = (X_1, \dots, X_d)^t \sim \text{mult}(n, p)$ , with  $p \in (0, 1)^d$  and  $\sum_{k=1}^d p_k = 1$

Under  $H_0 : p = (p_{0,1}, \dots, p_{0,d})^t$  it holds (approximately)

$$X^2 := \sum_{k=1}^d \frac{(X_k - \mathbb{E}_{H_0}[X_k])^2}{\mathbb{E}_{H_0}[X_k]} \stackrel{d}{\approx} \chi^2(d-1)$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2(d-1)$  as  $n \rightarrow \infty$

- here:  $n = 120$  and  $d = 6$ , as well as  $p_0 = (1/6, \dots, 1/6)^t$
- Let  $q_{1-\alpha}$  denote the  $(1 - \alpha)$ -quantile of the  $\chi^2(d-1)$ -distribution
- Here: can not reject  $H_0$





# The $\chi^2$ -test (goodness of fit)



- Let  $\mathfrak{X} = (X_1, \dots, X_d)^t \sim \text{mult}(n, p)$ , with  $p \in (0, 1)^d$  and  $\sum_{k=1}^d p_k = 1$

Under  $H_0 : p = (p_{0,1}, \dots, p_{0,d})^t$  it holds (approximately)

$$X^2 := \sum_{k=1}^d \frac{(X_k - \mathbb{E}_{H_0}[X_k])^2}{\mathbb{E}_{H_0}[X_k]} \stackrel{d}{\approx} \chi^2(d-1)$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2(d-1)$  as  $n \rightarrow \infty$


- here:  $n = 120$  and  $d = 6$ , as well as  $p_0 = (1/6, \dots, 1/6)^t$
- Let  $q_{1-\alpha}$  denote the  $(1 - \alpha)$ -quantile of the  $\chi^2(d-1)$ -distribution
- Here: can not reject  $H_0$



For details see the lecture: 'The  $\chi^2$ -test (goodness of fit)'

# The $\chi^2$ -test for independence

- Model: Let  $\mathfrak{X} = (X_{1,1}, \dots, X_{d_1, d_2})^t \sim \text{mult}(n, p)$   
with  $p = (p_{1,1}, \dots, p_{d_1, d_2})^t \in (0, 1)^{d_1 \cdot d_2}$  and  $\sum_{j,k} p_{j,k} = 1$



# The $\chi^2$ -test for independence

- Model: Let  $\mathfrak{X} = (X_{1,1}, \dots, X_{d_1,d_2})^t \sim \text{mult}(n, p)$   
with  $p = (p_{1,1}, \dots, p_{d_1,d_2})^t \in (0, 1)^{d_1 \cdot d_2}$  and  $\sum_{j,k} p_{j,k} = 1$

Under  $H_0 : p = p_0 := (p_{1,\cdot} \cdot p_{\cdot,1}, \dots, p_{d_1,\cdot} \cdot p_{\cdot,d_2})^t \in (0, 1)^{d_1 \cdot d_2}$



it holds (approximately)

$$\text{and } \sum_{j=1}^{d_1} p_{j,\cdot} = \sum_{k=1}^{d_2} p_{\cdot,k} = 1$$

$$X^2 := \sum_{j,k} \frac{\left( \underset{\text{red}}{X_{j,k}} - \frac{\underset{\text{red}}{X_{j,\cdot}} \cdot \underset{\text{red}}{X_{\cdot,k}}}{n} \right)^2}{\frac{\underset{\text{red}}{X_{j,\cdot}} \cdot \underset{\text{red}}{X_{\cdot,k}}}{n}} \stackrel{d}{\approx} \chi^2((d_1 - 1) \cdot (d_2 - 1))$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2((d_1 - 1)(d_2 - 1))$  as  $n \rightarrow \infty$

- Here:  $d_1 = 2, d_2 = 4$ , i.e.,  $X^2 \stackrel{H_0}{\sim} \chi^2(3)$  (approx)
- $\underset{\text{red}}{X_{j,\cdot}}, \underset{\text{red}}{X_{\cdot,k}}, \underset{\text{blue}}{p_{j,\cdot}}$  and  $\underset{\text{blue}}{p_{\cdot,k}}$  are the 'marginal frequencies / probabilities'



# The $\chi^2$ -test for independence

- Model: Let  $\mathfrak{X} = (X_{1,1}, \dots, X_{d_1, d_2})^t \sim \text{mult}(n, p)$   
with  $p = (p_{1,1}, \dots, p_{d_1, d_2})^t \in (0, 1)^{d_1 \cdot d_2}$  and  $\sum_{j,k} p_{j,k} = 1$

Under  $H_0 : p = p_0 := (p_{1,\cdot} \cdot p_{\cdot,1}, \dots, p_{d_1,\cdot} \cdot p_{\cdot,d_2})^t \in (0, 1)^{d_1 \cdot d_2}$



it holds (approximately)

$$\text{and } \sum_{j=1}^{d_1} p_{j,\cdot} = \sum_{k=1}^{d_2} p_{\cdot,k} = 1$$

$$X^2 := \sum_{j,k} \frac{\left( X_{j,k} - \frac{X_{j,\cdot} \cdot X_{\cdot,k}}{n} \right)^2}{\frac{X_{j,\cdot} \cdot X_{\cdot,k}}{n}} \stackrel{d}{\approx} \chi^2((d_1 - 1) \cdot (d_2 - 1))$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2((d_1 - 1)(d_2 - 1))$  as  $n \rightarrow \infty$

- Here:  $d_1 = 2, d_2 = 4$ , i.e.,  $X^2 \stackrel{H_0}{\sim} \chi^2(3)$  (approx)
- $X_{j,\cdot}$ ,  $X_{\cdot,k}$ ,  $p_{j,\cdot}$  and  $p_{\cdot,k}$  are the 'marginal frequencies / probabilities'
- Let  $q_{1-\alpha}$  denote the  $(1 - \alpha)$ -quantile of the  $\chi^2([d_1 - 1] \cdot [d_2 - 1])$ -distr.



# The $\chi^2$ -test for independence

- Model: Let  $\mathfrak{X} = (X_{1,1}, \dots, X_{d_1, d_2})^t \sim \text{mult}(n, p)$   
with  $p = (p_{1,1}, \dots, p_{d_1, d_2})^t \in (0, 1)^{d_1 \cdot d_2}$  and  $\sum_{j,k} p_{j,k} = 1$



Under  $H_0 : p = p_0 := (p_{1,\cdot} \cdot p_{\cdot,1}, \dots, p_{d_1,\cdot} \cdot p_{\cdot,d_2})^t \in (0, 1)^{d_1 \cdot d_2}$

it holds (approximately) and  $\sum_{j=1}^{d_1} p_{j,\cdot} = \sum_{k=1}^{d_2} p_{\cdot,k} = 1$

$$X^2 := \sum_{j,k} \frac{\left( X_{j,k} - \frac{X_{j,\cdot} \cdot X_{\cdot,k}}{n} \right)^2}{\frac{X_{j,\cdot} \cdot X_{\cdot,k}}{n}} \stackrel{d}{\approx} \chi^2((d_1 - 1) \cdot (d_2 - 1))$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2((d_1 - 1)(d_2 - 1))$  as  $n \rightarrow \infty$

- Here:  $d_1 = 2, d_2 = 4$ , i.e.,  $X^2 \stackrel{H_0}{\sim} \chi^2(3)$  (approx)
- $X_{j,\cdot}$ ,  $X_{\cdot,k}$ ,  $p_{j,\cdot}$  and  $p_{\cdot,k}$  are the 'marginal frequencies / probabilities'
- Let  $q_{1-\alpha}$  denote the  $(1 - \alpha)$ -quantile of the  $\chi^2([d_1 - 1] \cdot [d_2 - 1])$ -distr.
- Here: reject  $H_0$



# The $\chi^2$ -test for independence

- Model: Let  $\mathfrak{X} = (X_{1,1}, \dots, X_{d_1, d_2})^t \sim \text{mult}(n, p)$   
with  $p = (p_{1,1}, \dots, p_{d_1, d_2})^t \in (0, 1)^{d_1 \cdot d_2}$  and  $\sum_{j,k} p_{j,k} = 1$

Under  $H_0 : p = p_0 := (p_{1,\cdot} \cdot p_{\cdot,1}, \dots, p_{d_1,\cdot} \cdot p_{\cdot,d_2})^t \in (0, 1)^{d_1 \cdot d_2}$



it holds (approximately)

$$\text{and } \sum_{j=1}^{d_1} p_{j,\cdot} = \sum_{k=1}^{d_2} p_{\cdot,k} = 1$$

$$X^2 := \sum_{j,k} \frac{\left( X_{j,k} - \frac{X_{j,\cdot} \cdot X_{\cdot,k}}{n} \right)^2}{\frac{X_{j,\cdot} \cdot X_{\cdot,k}}{n}} \stackrel{d}{\approx} \chi^2((d_1 - 1) \cdot (d_2 - 1))$$

in fact, it holds that  $X^2 \xrightarrow{d} \chi^2((d_1 - 1)(d_2 - 1))$  as  $n \rightarrow \infty$

- Here:  $d_1 = 2, d_2 = 4$ , i.e.,  $X^2 \stackrel{H_0}{\sim} \chi^2(3)$  (approx)
- $X_{j,\cdot}$ ,  $X_{\cdot,k}$ ,  $p_{j,\cdot}$  and  $p_{\cdot,k}$  are the 'marginal frequencies / probabilities'
- Let  $q_{1-\alpha}$  denote the  $(1 - \alpha)$ -quantile of the  $\chi^2([d_1 - 1] \cdot [d_2 - 1])$ -distr.
- Here: reject  $H_0$



For details see the lecture: 'The  $\chi^2$ -test for independence'

Thank you!