


# Probabilistic Reasoning

## Chapter 14 (Models)



# Probabilistic Models

---

- Models describe how (a portion of) the world works

# Probabilistic Models

---

- Models describe how (a portion of) the world works
- **Models are always simplifications**
  - May not account for every variable
  - May not account for all interactions between variables
  - “All models are wrong; but some are useful.”
    - George E. P. Box

# Probabilistic Models

---

- Models describe how (a portion of) the world works
- Models are always simplifications
  - May not account for every variable
  - May not account for all interactions between variables
  - “All models are wrong; but some are useful.”
    - George E. P. Box
- What do we do with probabilistic models?
  - We (or our agents) need to reason about unknown variables, given evidence
  - Example: explanation (diagnostic reasoning)
  - Example: prediction (causal reasoning)
  - Example: value of information

# Bayes' Nets: Big Picture

---

- Two problems with using full joint distribution tables as our probabilistic models:
  - Unless there are only a few variables, the joint is WAY too big to represent explicitly. For  $n$  variables with domain size  $d$ , joint table has  $d^n$  entries --- exponential in  $n$ .

# Bayes' Nets: Big Picture

---

- Two problems with using full joint distribution tables as our probabilistic models:
  - Unless there are only a few variables, the joint is WAY too big to represent explicitly. For  $n$  variables with domain size  $d$ , joint table has  $d^n$  entries --- exponential in  $n$ .
  - Hard to learn (estimate) anything empirically about more than a few variables at a time

# Bayes' Nets: Big Picture

---

- Two problems with using full joint distribution tables as our probabilistic models:
  - Unless there are only a few variables, the joint is WAY too big to represent explicitly. For  $n$  variables with domain size  $d$ , joint table has  $d^n$  entries --- exponential in  $n$ .
  - Hard to learn (estimate) anything empirically about more than a few variables at a time
- **Bayes' nets**: a technique for describing complex joint distributions (models) using simple, local distributions (conditional probabilities)
  - More properly called **graphical models**
  - We describe how variables locally interact
  - Local interactions chain together to give global, indirect interactions

# Bayes' Nets

---

- Representation

- Informal first introduction of Bayes' nets through causality “intuition”
- More formal introduction of Bayes' nets



# Bayes' Nets

---

- Representation
  - Informal first introduction of Bayes' nets through causality “intuition”
  - More formal introduction of Bayes' nets
- Conditional Independences

# Bayes' Nets

---

- Representation
  - Informal first introduction of Bayes' nets through causality “intuition”
  - More formal introduction of Bayes' nets
- Conditional Independences
- Probabilistic Inference

# Bayes' Nets

---

- Representation
  - Informal first introduction of Bayes' nets through causality “intuition”
  - More formal introduction of Bayes' nets
- Conditional Independences
- Probabilistic Inference
- Learning Bayes' Nets from Data

# Graphical Model Notation

---

- Nodes: variables (with domains)
  - Can be assigned (observed) or unassigned (unobserved)



# Graphical Model Notation

---


- Nodes: variables (with domains)

- Can be assigned (observed) or unassigned (unobserved)



- Arcs: interactions


- Similar to CSP constraints
- Indicate “direct influence” between variables
- Formally: encode conditional independence (more later)



# Graphical Model Notation

---

- **Nodes: variables (with domains)**
  - Can be assigned (observed) or unassigned (unobserved)
- **Arcs: interactions**
  - Similar to CSP constraints
  - Indicate “direct influence” between variables
  - Formally: encode conditional independence (more later)
- **For now: imagine that arrows mean direct causation (in general, they don't!)**



# Example: Coin Flips

---

- N independent coin flips

$$X_1$$

$$X_2$$

...

$$X_n$$

# Example: Coin Flips

---

- N independent coin flips



- No interactions between variables:  
absolute independence



# Example: Traffic

---

- Variables:
  - R: It rains
  - T: There is traffic

# Example: Traffic


---

- Variables:
  - R: It rains
  - T: There is traffic
- Model 1: independence

# Example: Traffic

---


- Variables:
  - $R$ : It rains
  - $T$ : There is traffic
- Model 1: independence
- Model 2: rain causes traffic



# Example: Traffic

---

- Variables:
  - R: It rains
  - T: There is traffic
- Model 1: independence
- Model 2: rain causes traffic
- Why is an agent using model 2 better?



# Example: Traffic II

---

- Let's build a causal graphical model
- Variables
  - T: Traffic
  - R: It rains
  - L: Low pressure
  - D: Roof drips
  - B: Ballgame
  - C: Cavity

# Example: Alarm Network

---

- Variables
  - B: Burglary
  - A: Alarm goes off
  - M: Mary calls
  - J: John calls
  - E: Earthquake!

# Bayes' Net Semantics

---

- Let's formalize the semantics of a Bayes' net

# Bayes' Net Semantics

---


- Let's formalize the semantics of a Bayes' net
- A set of nodes, one per variable  $X$



# Bayes' Net Semantics

---

- Let's formalize the semantics of a Bayes' net
- A set of nodes, one per variable  $X$
- A directed, acyclic graph




# Bayes' Net Semantics

---

- Let's formalize the semantics of a Bayes' net
- A set of nodes, one per variable  $X$
- A directed, acyclic graph
- A conditional distribution for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$



$$P(X|A_1 \dots A_n)$$


# Bayes' Net Semantics

---

- Let's formalize the semantics of a Bayes' net
- A set of nodes, one per variable  $X$
- A directed, acyclic graph
- A conditional distribution for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values


$$P(X|a_1 \dots a_n)$$

- CPT: conditional probability table
- Description of a noisy “causal” process



# Probabilities in BNs


---



- Bayes' nets **implicitly** encode joint distributions
  - As a product of local conditional distributions

# Probabilities in BNs

---




- Bayes' nets **implicitly** encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

# Probabilities in BNs

---



- Bayes' nets **implicitly** encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:


$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

- Example:

$$P(+cavity, +catch, \neg toothache)$$

# Probabilities in BNs

---



- Bayes' nets **implicitly** encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:


$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

- Example:

$$P(+cavity, +catch, \neg toothache)$$

- This lets us reconstruct any entry of the full joint

# Probabilities in BNs



- Bayes' nets **implicitly** encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

- Example:


$$P(+cavity, +catch, \neg toothache)$$

- This lets us reconstruct any entry of the full joint
- Not every BN can represent every joint distribution**
  - The topology enforces certain conditional independencies



# Example: Coin Flips

---




$$P(h, h, t, h) =$$

*Only distributions whose variables are absolutely independent can be represented by a Bayes' net with no arcs.*

# Example: Traffic

---




$P(R)$

$+r$	$1/4$
$\neg r$	$3/4$

# Example: Traffic

---



$P(R)$

$+r$	$1/4$
$\neg r$	$3/4$


$P(T|R)$

$+r \rightarrow$

$+t$	$3/4$
$\neg t$	$1/4$

# Example: Traffic

---



$P(R)$

$+r$	$1/4$
$\neg r$	$3/4$


$P(T|R)$

$+r \rightarrow$	$+t$	$3/4$
	$\neg t$	$1/4$

$\neg r \rightarrow$	$+t$	$1/2$
	$\neg t$	$1/2$

# Example: Traffic

---



$P(R)$

$+r$	$1/4$
$\neg r$	$3/4$


$$P(+r, \neg t) =$$

$P(T|R)$

$+r \rightarrow$	$+t$	$3/4$
	$\neg t$	$1/4$
$\neg r \rightarrow$	$+t$	$1/2$
	$\neg t$	$1/2$

# Example: Alarm Network

B	P(B)
+b	0.001
¬b	0.999




E	P(E)
+e	0.002
¬e	0.998

A	J	P(J A)
+a	+j	0.9
+a	¬j	0.1
¬a	+j	0.05
¬a	¬j	0.95

A	M	P(M A)
+a	+m	0.7
+a	¬m	0.3
¬a	+m	0.01
¬a	¬m	0.99


B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	¬a	0.05
+b	¬e	+a	0.94
+b	¬e	¬a	0.06
¬b	+e	+a	0.29
¬b	+e	¬a	0.71
¬b	¬e	+a	0.001
¬b	¬e	¬a	0.999

# Example Bayes' Net: Insurance



# Example Bayes' Net: Car

---





# Build your own Bayes nets!

---

- <http://www.aispace.org/bayes/index.shtml>

# Size of a Bayes' Net

---

- How big is a joint distribution over  $N$  Boolean variables?

# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$

# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$
- How big is an N-node net if nodes have up to k parents?

# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$
- How big is an N-node net if nodes have up to k parents?  
 $O(N * 2^{k+1})$

# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$
- How big is an N-node net if nodes have up to k parents?  
 $O(N * 2^{k+1})$
- Both give you the power to calculate  $P(X_1, X_2, \dots, X_n)$

# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$
- How big is an N-node net if nodes have up to k parents?  
 $O(N * 2^{k+1})$
- Both give you the power to calculate  $P(X_1, X_2, \dots, X_n)$
- BNs: Huge space savings!

# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$
- How big is an N-node net if nodes have up to k parents?  
 $O(N * 2^{k+1})$
- Both give you the power to calculate  $P(X_1, X_2, \dots, X_n)$
- BNs: Huge space savings!
- Also easier to elicit local CPTs



# Size of a Bayes' Net

---

- How big is a joint distribution over N Boolean variables?  
 $2^N$
- How big is an N-node net if nodes have up to k parents?  
 $O(N * 2^{k+1})$
- Both give you the power to calculate  $P(X_1, X_2, \dots, X_n)$
- BNs: Huge space savings!
- Also easier to elicit local CPTs
- Also turns out to be faster to answer queries (coming)

# Bayes' Nets

---

- Representation

- ✓ Informal first introduction of Bayes' nets through causality “intuition”
  - More formal introduction of Bayes' nets

# Representing Joint Probability Distributions

---

- **Table representation:**

number of parameters:  $d^n - 1$

# Representing Joint Probability Distributions

---

- **Table representation:**

number of parameters:  $d^{n-1}$

- **Chain rule representation:**

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | x_1 \dots x_{i-1})$$

number of parameters:  $(d-1) + d(d-1) + d^2(d-1) + \dots + d^{n-1}(d-1) = d^n - 1$

Size of CPT = (number of different joint instantiations of the preceding variables)  
*times* (number of values current variable can take on *minus* 1)

# Representing Joint Probability Distributions

---

- **Table representation:**

number of parameters:  $d^n - 1$

- **Chain rule representation:**

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | x_1 \dots x_{i-1})$$

number of parameters:  $(d-1) + d(d-1) + d^2(d-1) + \dots + d^{n-1}(d-1) = d^n - 1$

Size of CPT = (number of different joint instantiations of the preceding variables)  
*times* (number of values current variable can take on *minus* 1)

- Both can represent any distribution over the  $n$  random variables.  
Makes sense same number of parameters needs to be stored.

# Representing Joint Probability Distributions

---

- **Table representation:**

number of parameters:  $d^{n-1}$

- **Chain rule representation:**

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | x_1 \dots x_{i-1})$$

number of parameters:  $(d-1) + d(d-1) + d^2(d-1) + \dots + d^{n-1}(d-1) = d^n - 1$

Size of CPT = (number of different joint instantiations of the preceding variables)  
*times* (number of values current variable can take on *minus* 1)

- Both can represent any distribution over the  $n$  random variables.  
Makes sense same number of parameters needs to be stored.
- Chain rule applies to all orderings of the variables, so for a given distribution we can represent it in  $n! = n \text{ factorial} = n(n-1)(n-2)\dots 2.1$  different ways with the chain rule

# Chain Rule $\rightarrow$ Bayes' net

- **Chain rule representation: applies to ALL distributions**

- Pick any ordering of variables, rename accordingly as  $x_1, x_2, \dots, x_n$

$$P(x_1, x_2, \dots, x_n) = \prod_i P(x_i | x_1 \dots x_{i-1})$$

**Exponential  
in n**

number of parameters:  $(d-1) + d(d-1) + d^2(d-1) + \dots + d^{n-1}(d-1) = d^n - 1$

# Chain Rule → Bayes' net

- **Chain rule representation: applies to ALL distributions**

- Pick any ordering of variables, rename accordingly as  $x_1, x_2, \dots, x_n$

$$P(x_1, x_2, \dots, x_n) = \prod_i P(x_i | x_1 \dots x_{i-1})$$

Exponential  
in  $n$

number of parameters:  $(d-1) + d(d-1) + d^2(d-1) + \dots + d^{n-1}(d-1) = d^n - 1$

- **Bayes' net representation: makes assumptions**

- Pick any ordering of variables, rename accordingly as  $x_1, x_2, \dots, x_n$
- Pick any directed acyclic graph consistent with the ordering
- Assume following conditional independencies:

$$P(x_i | x_1 \dots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

→ **Joint:**  $P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$

number of parameters: (maximum number of parents =  $K$ )

$$\sum_{i=1}^n d^{|\text{parents}(X_i)|} (d-1) = O(nd^K (d-1)) = O(nd^{K+1})$$

Linear  
in  $n$

Note: no causality assumption made anywhere.



# Causality?

---

- When Bayes' nets reflect the true causal patterns:
  - Often simpler (nodes have fewer parents)
  - Often easier to think about
  - Often easier to elicit from experts

# Causality?

---

- When Bayes' nets reflect the true causal patterns:
  - Often simpler (nodes have fewer parents)
  - Often easier to think about
  - Often easier to elicit from experts
- BNs need not actually be causal
  - Sometimes no causal net exists over the domain
  - E.g. consider the variables *Traffic* and *Drips*
  - End up with arrows that reflect correlation, not causation

# Causality?


---

- When Bayes' nets reflect the true causal patterns:
  - Often simpler (nodes have fewer parents)
  - Often easier to think about
  - Often easier to elicit from experts
- BNs need not actually be causal
  - Sometimes no causal net exists over the domain
  - E.g. consider the variables *Traffic* and *Drips*
  - End up with arrows that reflect correlation, not causation
- What do the arrows really mean?
  - Topology may happen to encode causal structure
  - **Topology only guaranteed to encode conditional independence**

# Example: Traffic


---

- Basic traffic net
- Let's multiply out the joint



# Example: Traffic

- Basic traffic net
- Let's multiply out the joint



$$P(R)$$

$r$	$1/4$
$\neg r$	$3/4$

$$P(T|R)$$

$r$	$t$	$3/4$
	$\neg t$	$1/4$

$\neg r$	$t$	$1/2$
	$\neg t$	$1/2$


$$P(T, R)$$

$r$	$t$	$3/16$
$r$	$\neg t$	$1/16$
$\neg r$	$t$	$6/16$
$\neg r$	$\neg t$	$6/16$

# Example: Reverse Traffic


---

- Reverse causality?



# Example: Reverse Traffic

- Reverse causality?



$$P(T)$$

t	9/16
$\neg t$	7/16

$$P(R|T)$$

t	r	1/3
	$\neg r$	2/3

$\neg t$	r	1/7
	$\neg r$	6/7



$$P(T, R)$$

r	t	3/16
r	$\neg t$	1/16
$\neg r$	t	6/16
$\neg r$	$\neg t$	6/16

# Example: Coins

---

- Extra arcs don't prevent representing independence, just allow non-independence



$P(X_1)$

h	0.5
t	0.5



$P(X_2)$

h	0.5
t	0.5



# Example: Coins

- Extra arcs don't prevent representing independence, just allow non-independence




$P(X_1)$

h	0.5
t	0.5

$P(X_2)$

h	0.5
t	0.5



$P(X_1)$

h	0.5
t	0.5


$P(X_2|X_1)$

h   h	0.5
t   h	0.5

h   t	0.5
t   t	0.5

# Example: Coins

- Extra arcs don't prevent representing independence, just allow non-independence



- Adding unneeded arcs isn't wrong, it's just inefficient

# Bayes' Nets

---

- Representation

- ✓ Informal first introduction of Bayes' nets through causality “intuition”
- ✓ More formal introduction of Bayes' nets

- Conditional Independences

- Probabilistic Inference

- Learning Bayes' Nets from Data

# Bayes Nets: Assumptions

---

- To go from chain rule to Bayes' net representation, we made the following assumption about the distribution:

$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

# Bayes Nets: Assumptions

---

- To go from chain rule to Bayes' net representation, we made the following assumption about the distribution:

$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Turns out that probability distributions that satisfy the above (“chain-rule→Bayes net”) conditional independence assumptions
  - often can be guaranteed to have many more conditional independences
  - These guaranteed additional conditional independences can be read off directly from the graph

# Bayes Nets: Assumptions

---


- To go from chain rule to Bayes' net representation, we made the following assumption about the distribution:

$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Turns out that probability distributions that satisfy the above (“chain-rule→Bayes net”) conditional independence assumptions
  - often can be guaranteed to have many more conditional independences
  - These guaranteed additional conditional independences can be read off directly from the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph

# Example


---



- Conditional independence assumptions directly from simplifications in chain rule:

# Example

---



- Conditional independence assumptions directly from simplifications in chain rule:
- Additional implied conditional independence assumptions?



# Independence in a BN

---

- Given a Bayes net graph

- Important question:

- Are two nodes guaranteed to be independent given certain evidence?*

# Independence in a BN

---

- Given a Bayes net graph

- Important question:

*Are two nodes guaranteed to be independent given certain evidence?*

Equivalent question:

*Are two nodes independent given the evidence in all distributions that can be encoded with the Bayes net graph?*

# Independence in a BN

---

- Given a Bayes net graph

- Important question:

*Are two nodes guaranteed to be independent given certain evidence?*

Equivalent question:

*Are two nodes independent given the evidence in all distributions that can be encoded with the Bayes net graph?*

- Before proceeding: How about opposite question: Are two nodes guaranteed to be *dependent* given certain evidence?

# Independence in a BN

---

- Given a Bayes net graph

- Important question:

*Are two nodes guaranteed to be independent given certain evidence?*

Equivalent question:

*Are two nodes independent given the evidence in all distributions that can be encoded with the Bayes net graph?*

- Before proceeding: How about opposite question: Are two nodes guaranteed to be *dependent* given certain evidence?

- No! For any BN graph you can choose all CPT's such that all variables are independent by having  $P(X \mid \text{Pa}(X)) = p_X$  not depend on the value of the parents. Simple way of doing so: pick all entries in all CPTs equal to 0.5 (assuming binary variables)

# D-separation: Outline

---

- Study independence properties for triples

# D-separation: Outline


---

- Study independence properties for triples
- Any complex example can be analyzed by considering relevant triples

# Causal Chains

---

- This configuration is a “causal chain”



$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

X: Low pressure


Y: Rain

Z: Traffic

# Causal Chains

---

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$


- Is it guaranteed that X is independent of Z ?



# Causal Chains

---

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Is it guaranteed that X is independent of Z ? **No!**

# Causal Chains

---

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic


$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Is it guaranteed that X is independent of Z ? **No!**
  - One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

# Causal Chains

---

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic


$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Is it guaranteed that X is independent of Z ? **No!**
  - One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.
  - Example:  $P(y|x) = 1$  if  $y=x$ , 0 otherwise  
 $P(z|y) = 1$  if  $z=y$ , 0 otherwise  
Then we have  $P(z|x) = 1$  if  $z=x$ , 0 otherwise  
hence X and Z are not independent in this example

# Causal Chains

---

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic


$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Is it guaranteed that X is independent of Z given Y?

$$P(z|x, y) = \frac{P(x, y, z)}{P(x, y)} = \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)}$$

# Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$


- Is it guaranteed that X is independent of Z given Y?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} = \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \end{aligned}$$

Yes!

# Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Is it guaranteed that X is independent of Z given Y?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} = \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \end{aligned}$$


Yes!

- Evidence along the chain “blocks” the influence

# Common Cause

---

- Another basic configuration: two effects of the same cause
  - Is it guaranteed that X and Z are independent?



Y: Project due


X: Piazza busy

Z: Lab full

# Common Cause

---

- Another basic configuration: two effects of the same cause
  - Is it guaranteed that X and Z are independent?
    - *No!*
    - Counterexample:  
Choose  $P(X|Y)=1$  if  $x=y$ , 0 otherwise,  
Choose  $P(z|y) = 1$  if  $z=y$ , 0 otherwise.



Y: Project due

X: Piazza busy

Z: Lab full



# Common Cause

---

- Another basic configuration: two effects of the same cause

- Is it guaranteed that  $X$  and  $Z$  are independent?


- *No!*

- Counterexample:

Choose  $P(X|Y)=1$  if  $x=y$ , 0 otherwise,

Choose  $P(z|y) = 1$  if  $z=y$ , 0 otherwise.

Then  $P(x|z)=1$  if  $x=z$  and 0 otherwise, hence  $X$  and  $Z$  are not independent in this example and hence it is not guaranteed that if a distribution can be encoded with the Bayes' net structure on the right that  $X$  and  $Z$  are independent in that distribution



Y: Project due

X: Piazza busy

Z: Lab full

# Common Cause


---

- Another basic configuration: two effects of the same cause

- Is it guaranteed that X and Z are independent given Y?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} = \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)} \\ &= P(z|y) \quad \text{Yes!} \end{aligned}$$

- Observing the cause blocks influence between effects.



Y: Project due


X: Piazza busy

Z: Lab full

# Common Effect

---

- Last configuration: two causes of one effect (v-structures)
  - Are X and Z independent?



X: Raining


Z: Ballgame

Y: Traffic

# Common Effect

---

- Last configuration: two causes of one effect (v-structures)
  - Are X and Z independent?
    - **Yes**: the ballgame and the rain cause traffic, but they are not correlated



X: Raining


Z: Ballgame

Y: Traffic

# Common Effect

---

- Last configuration: two causes of one effect (v-structures)
  - Are X and Z independent?
    - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
    - Still need to prove they must be (try it!)



X: Raining


Z: Ballgame

Y: Traffic

# Common Effect

---

- Last configuration: two causes of one effect (v-structures)
  - Are X and Z independent?
    - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
    - Still need to prove they must be (try it!)
  - Are X and Z independent given Y?



X: Raining


Z: Ballgame

Y: Traffic

# Common Effect

---

- Last configuration: two causes of one effect (v-structures)
  - Are X and Z independent?
    - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
    - Still need to prove they must be (try it!)
  - Are X and Z independent given Y?
    - **No**: seeing traffic puts the rain and the ballgame in competition as explanation?



X: Raining


Z: Ballgame

Y: Traffic

# Common Effect

---

- Last configuration: two causes of one effect (v-structures)
  - Are X and Z independent?
    - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
    - Still need to prove they must be (try it!)
  - Are X and Z independent given Y?
    - **No**: seeing traffic puts the rain and the ballgame in competition as explanation?
  - **This is backwards from the other cases**
    - Observing an effect **activates** influence between possible causes.



X: Raining

Z: Ballgame

Y: Traffic



# Reachability (D-Separation)

---


- Question: Are  $X$  and  $Y$  conditionally independent given evidence vars  $\{Z\}$ ?




# Reachability (D-Separation)

- Question: Are  $X$  and  $Y$  conditionally independent given evidence vars  $\{Z\}$ ?
  - Consider all (undirected) paths from  $X$  to  $Y$
  - No active paths = independence!

Active Triples





Inactive Triples





# Reachability (D-Separation)

- Question: Are X and Y conditionally independent given evidence vars {Z}?
- Consider all (undirected) paths from X to Y
- No active paths = independence!
- A path is active if each triple is active:
  - Causal chain  $A \rightarrow B \rightarrow C$  where B is unobserved (either direction)
  - Common cause  $A \leftarrow B \rightarrow C$  where B is unobserved

Active Triples






Inactive Triples





# Reachability (D-Separation)

- Question: Are X and Y conditionally independent given evidence vars {Z}?
- Consider all (undirected) paths from X to Y
- No active paths = independence!
- A path is active if each triple is active:
  - Causal chain  $A \rightarrow B \rightarrow C$  where B is unobserved (either direction)
  - Common cause  $A \leftarrow B \rightarrow C$  where B is unobserved
  - Common effect (aka v-structure)  $A \rightarrow B \leftarrow C$  where B or one of its descendants is observed

Active Triples



Inactive Triples



# D-Separation

---

- Given query  $X_i \stackrel{?}{\perp} X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$

# D-Separation

---

- Given query  $X_i \overset{?}{\perp} X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$
- Shade all evidence nodes

# D-Separation

---

- Given query  $X_i \overset{?}{\perp\!\!\!\perp} X_j | \{X_{k_1}, \dots, X_{k_n}\}$
- Shade all evidence nodes
- For all (undirected!) paths between and
  - Check whether path is active
    - If active return:  
not guaranteed that  $X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$

# D-Separation

---

- Given query  $X_i \stackrel{?}{\perp\!\!\!\perp} X_j | \{X_{k_1}, \dots, X_{k_n}\}$
- Shade all evidence nodes
- For all (undirected!) paths between and
  - Check whether path is active
    - If active return:  
not guaranteed that  $X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$
- (If reaching this point all paths have been checked and shown inactive)
  - Return: guaranteed that  $X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$




# Example

---

$$R \perp\!\!\!\perp B$$

$$R \perp\!\!\!\perp B | T$$

$$R \perp\!\!\!\perp B | T'$$



# Example


---

$$R \perp\!\!\!\perp B$$

Yes

$$R \perp\!\!\!\perp B | T$$

$$R \perp\!\!\!\perp B | T'$$



# Example

---


$$L \perp\!\!\!\perp T' | T$$

$$L \perp\!\!\!\perp B$$

$$L \perp\!\!\!\perp B | T$$

$$L \perp\!\!\!\perp B | T'$$

$$L \perp\!\!\!\perp B | T, R$$



# Example

---


$L \perp\!\!\!\perp T' | T$       Yes

$L \perp\!\!\!\perp B$       Yes

$L \perp\!\!\!\perp B | T$

$L \perp\!\!\!\perp B | T'$

$L \perp\!\!\!\perp B | T, R$       Yes




# Example

---

- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad
- Questions:

$$T \perp\!\!\!\perp D$$




# Example

---

- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad
- Questions:

$$T \perp\!\!\!\perp D$$


$$T \perp\!\!\!\perp D | R$$



# Example

---

- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad



- Questions:

$$T \perp\!\!\!\perp D$$


$$T \perp\!\!\!\perp D | R$$

Yes

# Example

---

- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad



- Questions:

$$T \perp\!\!\!\perp D$$

$$T \perp\!\!\!\perp D | R$$

Yes

$$T \perp\!\!\!\perp D | R, S$$



# All Conditional Independences

---

- Given a Bayes net structure, can run d-separation to build a complete list of conditional independences that are guaranteed to be true, all of the form

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$


# Topology Limits Distributions

---

- Given some graph topology  $G$ , only certain joint distributions can be encoded


# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences




# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)




# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs



# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution



# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions

# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure



# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone

# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

# Bayes' Nets

---



Representation



Conditional Independences

- Probabilistic Inference

- Enumeration (exact, exponential complexity)
- Variable elimination (exact, worst-case exponential complexity, often better)
- Probabilistic inference is NP-complete
- Sampling (approximate)

- Learning Bayes' Nets from Data

\_\_\_\_\_