

Quantifying Uncertainty

Chapter 13

Acting Under Uncertainty

Partial sensor info is badly handled in logic because

- Every possible explanation has to be accounted for
 - No matter how unlikely the explanation is
 - Which leads to very large and complex belief-states (BS) where
 - BS are a representation of the set of all possible world states
- Every explanation must have a contingency action
 - No matter how unlikely the contingency is
 - Hence correct contingency plans can grow arbitrarily large
- Often there is no plan guaranteed to achieve the goal
 - Yet the agent must act and therefore it
 - Must have a way of comparing the merits of alternative plans

Acting Under Uncertainty

Taxi driver: Deliver a passenger on time at the airport

- **Form a plan A90:** meaning that it
 - Leaves home 90 minutes before the flight departs and
 - Driving at a reasonable speed to not get a ticket
- **Even though the airport is only 5' away** a logical TD agent
 - Will not be able to conclude that plan A90 is true. Instead it says
 - A90 is true as long as: car doesn't break, run out of gas, meteorite ...
- **None of these conditions** can be deduced for sure, so
 - The plan's success cannot be inferred
 - This is called the qualification problem!

Acting Under Uncertainty

Taxi driver: Statistical approach

- **Plan A90:** does in some sense the right thing as
 - It is expected to maximize the agent's performance measure (PM)
 - PM: arriving on time, avoiding tickets, and long wait at airport
- **Agent's knowledge** cannot guarantee with certainty A90
 - But it can provide some degree of belief that it can be achieved
 - A120 will increase certainty, but also the likelihood of a long wait
- **The rational decision** depends on the relative importance
 - Of the various goals such as arriving on time, no speeding, wait time
 - And the degree to which they will be achieved

Summarizing Uncertainty

Diagnosing a dental patient's toothache (uncertainty)

- Consider the rule: $\text{Toothache} \rightarrow \text{Cavity}$
 - Problem: this rule is wrong!
 - Not all patients with toothaches have cavities.
- Try fix 1: $\text{Toothache} \rightarrow \text{Cavity} \vee \text{Abscess} \vee \text{GumDisease} \vee \dots$
 - Problem: to make rule true, one has to add an unlimited list
- Try fix 2: $\text{Cavity} \rightarrow \text{Toothache}$
 - Problem: this rule is wrong!
 - Not all cavities cause pain

Summarizing Uncertainty

Logic fails in medicine for three main reasons:

- **Laziness:** It is too much work to
 - List the complete set of antecedents or consequents
 - Such rules are too hard to use if these lists are complete
- **Ignorance 1:** Medicine has no complete theory
 - Hence, there is no way to write complete antecedents or consequents
- **Ignorance 2:** Even if we know all the rules
 - We may be uncertain about a particular patient because
 - Not all necessary tests have been or can be run

Other similar domains: law, business, repair, dating,...

Summarizing Uncertainty

Agent's knowledge can only provide a degree of belief

- **Probability theory:** The main tool for handling such degrees

Ontological commitments of logic and probability theory

- **The same:** The word consists of facts that hold or do not hold

Epistemological commitments of logic and probability theory

- **Logic:** believes such facts to be 0, 1, or no opinion
- **Probability theory:** has a numerical degree of belief between 0 and 1

Probability theory allows to summarize uncertainty

- **With source:** being laziness and ignorance
- **PT statements:** reflect a knowledge state, and not a word state

Uncertainty and Rational Decisions

Is A180 better than A120 or A90?

- **Preferences:** Allow a rational agent to decide which one is better
- **Outcome:** A completely specified state, including preference values


Utility theory allows to represent and reason with preferences

- **Utility:** Every state has a degree of usefulness or utility to the agent
- **Preference:** The agent will prefer states with higher utility

Decision theory combines probabilities with utilities

- **Decision Theory** = Probability Theory + Utility Theory
- **Rational** = Chose actions that maximize the expected utility (MEU)

Decision-Theoretic Agent



function Decision-Theoretic-Agent (**percept**) **returns** **action**
persistent state, action, model, rules

state = Update-State(state, action, percept, model)

probabilities = compute outcome probabilities for all actions given state and model

action = select action with highest expected utility given probabilities and utility info

return action

Probabilistic Reasoning

- Probability

Probabilistic Reasoning

- Probability
- Distributions over LARGE Numbers of Random Variables

Probabilistic Reasoning

- Probability
- Distributions over LARGE Numbers of Random Variables
 - Representation

Probabilistic Reasoning

- Probability
- Distributions over LARGE Numbers of Random Variables
 - Representation
 - Independence

Probabilistic Reasoning

- Probability
- Distributions over LARGE Numbers of Random Variables
 - Representation
 - Independence
 - Inference
 - Variable Elimination
 - Sampling

Probabilistic Reasoning

- Probability
- Distributions over LARGE Numbers of Random Variables
 - Representation
 - Independence
 - Inference
 - Variable Elimination
 - Sampling
 - Hidden Markov Models

Random Variables

- A random variable is some aspect of the world about which we (may) have uncertainty
 - R = Is it raining?
 - D = How long will it take to drive to work?
 - L = Where am I?

Random Variables

- A random variable is some aspect of the world about which we (may) have uncertainty
 - R = Is it raining?
 - D = How long will it take to drive to work?
 - L = Where am I?
- We denote random variables with capital letters

Random Variables

- A random variable is some aspect of the world about which we (may) have uncertainty
 - R = Is it raining?
 - D = How long will it take to drive to work?
 - L = Where am I?
- We denote random variables with capital letters
- Like variables in a CSP, random variables have domains
 - R in $\{\text{true}, \text{false}\}$ (sometimes write as $\{+r, -r\}$)
 - D in $[0, \infty)$
 - L in possible locations, maybe $\{(0,0), (0,1), \dots\}$

Probability Distributions

- Random variables have distributions

$P(T)$

T	P
warm	0.5
cold	0.5

$P(W)$

W	P
sun	0.6
rain	0.1
fog	0.3
meteor	0.0

Probability Distributions

- Random variables have distributions

$P(T)$		$P(W)$	
T	P	W	P
warm	0.5	sun	0.6
cold	0.5	rain	0.1
		fog	0.3
		meteor	0.0

- A distribution is a TABLE of probabilities of values

Probability Distributions

- Random variables have distributions

$P(T)$		$P(W)$	
T	P	W	P
warm	0.5	sun	0.6
cold	0.5	rain	0.1
		fog	0.3
		meteor	0.0

- A distribution is a TABLE of probabilities of values
- A probability (lower case value) is a single number

$$P(W = \text{rain}) = 0.1$$

$$P(\text{rain}) = 0.1$$

Probability Distributions

- Random variables have distributions

$P(T)$		$P(W)$	
T	P	W	P
warm	0.5	sun	0.6
cold	0.5	rain	0.1
		fog	0.3
		meteor	0.0

- A distribution is a TABLE of probabilities of values
- A probability (lower case value) is a single number

$$P(W = \text{rain}) = 0.1$$

$$P(\text{rain}) = 0.1$$

- Must have:** $\forall x P(x) \geq 0$ $\sum_x P(x) = 1$

Joint Distributions

- A *joint distribution* over a set of random variables: X_1, X_2, \dots, X_n specifies a real number for each assignment (or *outcome*):

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

$$P(x_1, x_2, \dots, x_n)$$

Joint Distributions

- A *joint distribution* over a set of random variables: X_1, X_2, \dots, X_n specifies a real number for each assignment (or *outcome*):

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

$$P(x_1, x_2, \dots, x_n)$$

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Joint Distributions

- A *joint distribution* over a set of random variables: X_1, X_2, \dots, X_n specifies a real number for each assignment (or *outcome*):

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

$$P(x_1, x_2, \dots, x_n)$$

- Size of distribution if n variables with domain sizes d?

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Joint Distributions

- A *joint distribution* over a set of random variables: X_1, X_2, \dots, X_n specifies a real number for each assignment (or *outcome*):

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

$$P(x_1, x_2, \dots, x_n)$$

- Size of distribution if n variables with domain sizes d ?
- Must obey:

$$P(x_1, x_2, \dots, x_n) \geq 0$$

$$\sum_{(x_1, x_2, \dots, x_n)} P(x_1, x_2, \dots, x_n) = 1$$

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Joint Distributions

- A *joint distribution* over a set of random variables: X_1, X_2, \dots, X_n specifies a real number for each assignment (or *outcome*):

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

$$P(x_1, x_2, \dots, x_n)$$

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

- Size of distribution if n variables with domain sizes d ?
- Must obey:

$$P(x_1, x_2, \dots, x_n) \geq 0$$

$$\sum_{(x_1, x_2, \dots, x_n)} P(x_1, x_2, \dots, x_n) = 1$$

- For all but the smallest distributions, impractical to write out

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*
 - Joint distributions: say whether assignments (outcomes) are likely

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*
 - Joint distributions: say whether assignments (outcomes) are likely
 - *Normalized*: sum to 1.0

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*
 - Joint distributions: say whether assignments (outcomes) are likely
 - *Normalized*: sum to 1.0
 - Ideally: only certain variables directly interact

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*
 - Joint distributions: say whether assignments (outcomes) are likely
 - *Normalized*: sum to 1.0
 - Ideally: only certain variables directly interact
- Constraint satisfaction probs:
 - Variables with domains

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Constraint over T,W

T	W	P
hot	sun	T
hot	rain	F
cold	sun	F
cold	rain	T

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*
 - Joint distributions: say whether assignments (outcomes) are likely
 - *Normalized*: sum to 1.0
 - Ideally: only certain variables directly interact
- Constraint satisfaction probs:
 - Variables with domains
 - Constraints: state whether assignments are possible

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Constraint over T,W

T	W	P
hot	sun	T
hot	rain	F
cold	sun	F
cold	rain	T

Probabilistic Models

- A probabilistic model is a joint distribution over a set of random variables
- Probabilistic models:
 - (Random) variables with domains
Assignments are called *outcomes*
 - Joint distributions: say whether assignments (outcomes) are likely
 - *Normalized*: sum to 1.0
 - Ideally: only certain variables directly interact
- Constraint satisfaction probs:
 - Variables with domains
 - Constraints: state whether assignments are possible
 - Ideally: only certain variables directly interact

Distribution over T,W

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Constraint over T,W

T	W	P
hot	sun	T
hot	rain	F
cold	sun	F
cold	rain	T

Events

- An *event* is a set E of outcomes

$$P(E) = \sum_{(x_1 \dots x_n) \in E} P(x_1 \dots x_n)$$

Events

- An *event* is a set E of outcomes

$$P(E) = \sum_{(x_1 \dots x_n) \in E} P(x_1 \dots x_n)$$

- From a joint distribution, we can calculate the probability of any event
 - Probability that it's hot AND sunny?

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Events

- An *event* is a set E of outcomes

$$P(E) = \sum_{(x_1 \dots x_n) \in E} P(x_1 \dots x_n)$$

- From a joint distribution, we can calculate the probability of any event
 - Probability that it's hot AND sunny?
 - Probability that it's hot?

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Events

- An *event* is a set E of outcomes

$$P(E) = \sum_{(x_1 \dots x_n) \in E} P(x_1 \dots x_n)$$

- From a joint distribution, we can calculate the probability of any event
 - Probability that it's hot AND sunny?
 - Probability that it's hot?
 - Probability that it's hot OR sunny?

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Events

- An *event* is a set E of outcomes

$$P(E) = \sum_{(x_1 \dots x_n) \in E} P(x_1 \dots x_n)$$

- From a joint distribution, we can calculate the probability of any event
 - Probability that it's hot AND sunny?
 - Probability that it's hot?
 - Probability that it's hot OR sunny?
- Typically, the events we care about are *partial assignments*, like $P(T=\text{hot})$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Marginal Distributions

- Marginal distributions are sub-tables which eliminate variables

Marginal Distributions

- Marginal distributions are sub-tables which eliminate variables
- Marginalization (summing out): Combine collapsed rows by adding

$P(T, W)$					$P(T)$	
T	W	P			T	P
hot	sun	0.4	$\xrightarrow{P(t) = \sum_s P(t, s)}$		hot	0.5
hot	rain	0.1			cold	0.5
cold	sun	0.2				
cold	rain	0.3				

Marginal Distributions

- Marginal distributions are sub-tables which eliminate variables
- Marginalization (summing out): Combine collapsed rows by adding

$P(T, W)$			$P(T)$	
T	W	P	T	P
hot	sun	0.4	hot	0.5
hot	rain	0.1	cold	0.5
cold	sun	0.2		
cold	rain	0.3		

$P(t) = \sum_s P(t, s)$

$P(W)$	
W	P
sun	0.6
rain	0.4

$P(s) = \sum_t P(t, s)$

Marginal Distributions

- Marginal distributions are sub-tables which eliminate variables
- Marginalization (summing out): Combine collapsed rows by adding

$P(T, W)$			$P(T)$	
T	W	P		
hot	sun	0.4	hot	0.5
hot	rain	0.1	cold	0.5
cold	sun	0.2	$P(W)$	
cold	rain	0.3	W	P

$P(t) = \sum_s P(t, s)$		sun	0.6
$P(s) = \sum_t P(t, s)$		rain	0.4

$$P(X_1 = x_1) = \sum_{x_2} P(X_1 = x_1, X_2 = x_2)$$

Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

$$(b \Leftarrow a) \wedge a$$

Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

$$a \wedge b = (b \Leftarrow a) \wedge a$$

Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

$$a \wedge b = (b \Leftarrow a) \wedge a$$


$$P(a, b) = P(b \mid a) P(a)$$

Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

$$a \wedge b = (b \Leftarrow a) \wedge a$$

$$P(a, b) = P(b | a) P(a)$$




Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

$$a \wedge b = (b \Leftarrow a) \wedge a = (a \Leftarrow b) \wedge b$$

$$P(a, b) = P(b | a) P(a) = P(a | b) P(b)$$




Conditional Probabilities

- A simple relation between joint and conditional probabilities
 - In fact, this is taken as the *definition* of a conditional probability

$$a \wedge b = (b \Leftarrow a) \wedge a = (a \Leftarrow b) \wedge b$$

$$P(a, b) = P(b | a) P(a) = P(a | b) P(b)$$



$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

$$P(W = r | T = c) = ???$$

Conditional Distributions

- Conditional distributions are probability distributions over some variables given fixed values of others

Conditional Distributions

$P(W T)$	$P(W T = \text{hot})$	
	W	P
	sun	0.8
	rain	0.2
	$P(W T = \text{cold})$	
	W	P
	sun	0.4
	rain	0.6

Joint Distribution

$P(T, W)$		
T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

Normalization Trick

- A trick to get a whole conditional distribution at once:
 - Select the joint probabilities matching the evidence
 - Normalize the selection (make it sum to one)

$$P(T, W)$$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

→
Select

$$P(T, r)$$

T	R	P
hot	rain	0.1
cold	rain	0.3

→
Normalize

$$P(T|r)$$

T	P
hot	0.25
cold	0.75

Normalization Trick

- A trick to get a whole conditional distribution at once:
 - Select the joint probabilities matching the evidence
 - Normalize the selection (make it sum to one)

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3



Select

$P(T, r)$

T	R	P
hot	rain	0.1
cold	rain	0.3



Normalize

$P(T|r)$

T	P
hot	0.25
cold	0.75

- Why does this work? Sum of selection is $P(\text{evidence})$! ($P(r)$, here)

$$P(x_1|x_2) = \frac{P(x_1, x_2)}{P(x_2)} = \frac{P(x_1, x_2)}{\sum_{x_1} P(x_1, x_2)}$$

Probabilistic Inference

- Probabilistic inference: compute a desired probability from other known probabilities (e.g. conditional from joint)

Probabilistic Inference

- Probabilistic inference: compute a desired probability from other known probabilities (e.g. conditional from joint)
- We generally compute conditional probabilities
 - $P(\text{on time} \mid \text{no reported accidents}) = 0.90$
 - These represent the agent's *beliefs* given the evidence

Probabilistic Inference

- Probabilistic inference: compute a desired probability from other known probabilities (e.g. conditional from joint)
- We generally compute conditional probabilities
 - $P(\text{on time} \mid \text{no reported accidents}) = 0.90$
 - These represent the agent's *beliefs* given the evidence
- Probabilities change with new evidence:
 - $P(\text{on time} \mid \text{no accidents, 5 a.m.}) = 0.95$
 - $P(\text{on time} \mid \text{no accidents, 5 a.m., raining}) = 0.80$
 - Observing new evidence causes *beliefs to be updated*

Inference by Enumeration

- $P(\text{sun})?$
- $P(\text{sun} \mid \text{winter})?$
- $P(\text{sun} \mid \text{winter, warm})?$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

Inference by Enumeration

- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- } X_1, X_2, \dots, X_n
All variables

Inference by Enumeration

- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- } X_1, X_2, \dots, X_n
All variables

- We want: $P(Q|e_1 \dots e_k)$

Inference by Enumeration

- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- $\left. \begin{array}{l} X_1, X_2, \dots X_n \\ \text{All variables} \end{array} \right\}$

- We want: $P(Q|e_1 \dots e_k)$

- First, select the entries consistent with the evidence
- Second, sum out H to get joint of Query and evidence:

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} \underbrace{P(Q, h_1 \dots h_r, e_1 \dots e_k)}_{X_1, X_2, \dots X_n}$$

Inference by Enumeration

- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- $\left. \begin{array}{l} X_1, X_2, \dots, X_n \\ \text{All variables} \end{array} \right\}$

- We want: $P(Q|e_1 \dots e_k)$

- First, select the entries consistent with the evidence
- Second, sum out H to get joint of Query and evidence:

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} \underbrace{P(Q, h_1 \dots h_r, e_1 \dots e_k)}_{X_1, X_2, \dots, X_n}$$

- Finally, normalize the remaining entries to conditionalize

Inference by Enumeration

- General case:

- Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 - Query* variable: Q
 - Hidden variables: $H_1 \dots H_r$
- $\left. \begin{array}{l} X_1, X_2, \dots X_n \\ \text{All variables} \end{array} \right\}$

- We want: $P(Q|e_1 \dots e_k)$

- First, select the entries consistent with the evidence
- Second, sum out H to get joint of Query and evidence:

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} \underbrace{P(Q, h_1 \dots h_r, e_1 \dots e_k)}_{X_1, X_2, \dots X_n}$$

- Finally, normalize the remaining entries to conditionalize
- Obvious problems:
 - Worst-case time complexity $O(d^n)$
 - Space complexity $O(d^n)$ to store the joint distribution

** Works fine with multiple query variables, too*

The Product Rule

- Sometimes have conditional distributions but want the joint

$$P(x|y) = \frac{P(x, y)}{P(y)} \quad \longleftrightarrow \quad P(x, y) = P(x|y)P(y)$$

The Product Rule

- Sometimes have conditional distributions but want the joint

$$P(x|y) = \frac{P(x, y)}{P(y)} \quad \longleftrightarrow \quad P(x, y) = P(x|y)P(y)$$

- Example:

$P(W)$

R	P
sun	0.8
rain	0.2

$P(D|W)$

D	W	P
wet	sun	0.1
dry	sun	0.9
wet	rain	0.7
dry	rain	0.3

$P(D, W)$

D	W	P
wet	sun	0.08
dry	sun	0.72
wet	rain	0.14
dry	rain	0.06

The Chain Rule

- More generally, can always write any joint distribution as an incremental product of conditional distributions

$$P(x_1, x_2, x_3) = P(x_1)P(x_2|x_1)P(x_3|x_1, x_2)$$

The Chain Rule

- More generally, can always write any joint distribution as an incremental product of conditional distributions

$$P(x_1, x_2, x_3) = P(x_1)P(x_2|x_1)P(x_3|x_1, x_2)$$

$$P(x_1, x_2, \dots x_n) = \prod_i P(x_i|x_1 \dots x_{i-1})$$

The Chain Rule

- More generally, can always write any joint distribution as an incremental product of conditional distributions

$$P(x_1, x_2, x_3) = P(x_1)P(x_2|x_1)P(x_3|x_1, x_2)$$

$$P(x_1, x_2, \dots x_n) = \prod_i P(x_i|x_1 \dots x_{i-1})$$

- Why is this always true?

The Chain Rule

- More generally, can always write any joint distribution as an incremental product of conditional distributions

$$P(x_1, x_2, x_3) = P(x_1)P(x_2|x_1)P(x_3|x_1, x_2)$$

$$P(x_1, x_2, \dots x_n) = \prod_i P(x_i|x_1 \dots x_{i-1})$$

- Why is this always true?
- Can now build a joint distributions only specifying conditionals!
 - Bayesian networks essentially apply the chain rule plus make conditional independence assumptions.

Bayes' Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

Bayes' Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

That's my rule!

- Dividing, we get:

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$



Bayes' Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

That's my rule!

- Dividing, we get:

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$

- Why is this at all helpful?

- Lets us build one conditional from its reverse
- Often one conditional is tricky but the other one is simple
- Foundation of many systems we'll see later (e.g. ASR, MT)



Bayes' Rule

- Two ways to factor a joint distribution over two variables:

$$P(x, y) = P(x|y)P(y) = P(y|x)P(x)$$

That's my rule!

- Dividing, we get:

$$P(x|y) = \frac{P(y|x)P(x)}{P(y)}$$

- Why is this at all helpful?
 - Lets us build one conditional from its reverse
 - Often one conditional is tricky but the other one is simple
 - Foundation of many systems we'll see later (e.g. ASR, MT)
- In the running for most important AI equation!



Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{Cause}|\text{Effect}) = \frac{P(\text{Effect}|\text{Cause})P(\text{Cause})}{P(\text{Effect})}$$

Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{Cause}|\text{Effect}) = \frac{P(\text{Effect}|\text{Cause})P(\text{Cause})}{P(\text{Effect})}$$

- Example:

- m is meningitis, s is stiff neck

$$P(s|m) = 0.8$$

$$P(m) = 0.0001$$

$$P(s) = 0.1$$

} Example
givens

Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{Cause}|\text{Effect}) = \frac{P(\text{Effect}|\text{Cause})P(\text{Cause})}{P(\text{Effect})}$$

- Example:

- m is meningitis, s is stiff neck

$$P(s|m) = 0.8$$

$$P(m) = 0.0001$$

$$P(s) = 0.1$$

} Example
givens

$$P(m|s) = \frac{P(s|m)P(m)}{P(s)} = \frac{0.8 \times 0.0001}{0.1} = 0.0008$$

Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{Cause}|\text{Effect}) = \frac{P(\text{Effect}|\text{Cause})P(\text{Cause})}{P(\text{Effect})}$$

- Example:

- m is meningitis, s is stiff neck

$$P(s|m) = 0.8$$

$$P(m) = 0.0001$$

$$P(s) = 0.1$$

} Example
gives

$$P(m|s) = \frac{P(s|m)P(m)}{P(s)} = \frac{0.8 \times 0.0001}{0.1} = 0.0008$$

- Note: posterior probability of meningitis still very small

Inference with Bayes' Rule

- Example: Diagnostic probability from causal probability:

$$P(\text{Cause}|\text{Effect}) = \frac{P(\text{Effect}|\text{Cause})P(\text{Cause})}{P(\text{Effect})}$$

- Example:

- m is meningitis, s is stiff neck

$$P(s|m) = 0.8$$

$$P(m) = 0.0001$$

$$P(s) = 0.1$$

} Example
gives

$$P(m|s) = \frac{P(s|m)P(m)}{P(s)} = \frac{0.8 \times 0.0001}{0.1} = 0.0008$$

- Note: posterior probability of meningitis still very small
 - Note: you should still get stiff necks checked out! Why?

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!
- Equivalent definition of independence:

$$\forall x, y : P(x|y) = P(x)$$

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!
- Equivalent definition of independence:

$$\forall x, y : P(x|y) = P(x)$$

- We write: $X \perp\!\!\!\perp Y$

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!
- Equivalent definition of independence:

$$\forall x, y : P(x|y) = P(x)$$

- We write: $X \perp\!\!\!\perp Y$

- Independence is a simplifying *modeling assumption*

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!
- Equivalent definition of independence:

$$\forall x, y : P(x|y) = P(x)$$

- We write: $X \perp\!\!\!\perp Y$

- Independence is a simplifying *modeling assumption*
 - Empirical* joint distributions: at best “close” to independent

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!
- Equivalent definition of independence:

$$\forall x, y : P(x|y) = P(x)$$

- We write: $X \perp\!\!\!\perp Y$

- Independence is a simplifying *modeling assumption*

- Empirical* joint distributions: at best “close” to independent
- What could we assume for {Weather, Traffic, Cavity, Toothache}?

Independence

- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- Says their joint distribution *factors* into a product two simpler ones.
- Usually variables are not independent!
- Equivalent definition of independence:

$$\forall x, y : P(x|y) = P(x)$$

- We write: $X \perp\!\!\!\perp Y$

- Independence is a simplifying *modeling assumption*
 - Empirical* joint distributions: at best “close” to independent
 - What could we assume for {Weather, Traffic, Cavity, Toothache}?
- Independence is like something from CSPs, what?

Example: Independence?

$P_1(T, W)$

T	W	P
warm	sun	0.4
warm	rain	0.1
cold	sun	0.2
cold	rain	0.3

$P(T)$

T	P
warm	0.5
cold	0.5

$P(W)$

W	P
sun	0.6
rain	0.4

Example: Independence?

$P_1(T, W)$

T	W	P
warm	sun	0.4
warm	rain	0.1
cold	sun	0.2
cold	rain	0.3

$P(T)$

T	P
warm	0.5
cold	0.5

$P_2(T, W)$


T	W	P
warm	sun	0.3
warm	rain	0.2
cold	sun	0.3
cold	rain	0.2

$P(W)$

W	P
sun	0.6
rain	0.4

Example: Independence

- N fair, independent coin flips:



Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$

Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$

Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$
- The same independence holds if I don't have a cavity:
 - $P(+\text{catch} \mid +\text{toothache}, \neg\text{cavity}) = P(+\text{catch} \mid \neg\text{cavity})$

Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$
- The same independence holds if I don't have a cavity:
 - $P(+\text{catch} \mid +\text{toothache}, \neg\text{cavity}) = P(+\text{catch} \mid \neg\text{cavity})$
- Catch is *conditionally independent* of Toothache given Cavity:
 - $P(\text{Catch} \mid \text{Toothache}, \text{Cavity}) = P(\text{Catch} \mid \text{Cavity})$

Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$
- The same independence holds if I don't have a cavity:
 - $P(+\text{catch} \mid +\text{toothache}, \neg\text{cavity}) = P(+\text{catch} \mid \neg\text{cavity})$
- Catch is *conditionally independent* of Toothache given Cavity:
 - $P(\text{Catch} \mid \text{Toothache}, \text{Cavity}) = P(\text{Catch} \mid \text{Cavity})$
- Equivalent statements:
 - $P(\text{Toothache} \mid \text{Catch}, \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity})$
 - $P(\text{Toothache}, \text{Catch} \mid \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity}) P(\text{Catch} \mid \text{Cavity})$
 - One can be derived from the other easily

Conditional Independence

- Unconditional (absolute) independence very rare (why?)

Conditional Independence

- Unconditional (absolute) independence very rare (why?)
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

$$\forall x, y, z : P(x|z, y) = P(x|z)$$

$$X \perp\!\!\!\perp Y | Z$$

Conditional Independence

- Unconditional (absolute) independence very rare (why?)
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

$$\forall x, y, z : P(x|z, y) = P(x|z)$$

$$X \perp\!\!\!\perp Y | Z$$

- What about this domain:
 - Traffic
 - Umbrella
 - Raining

Conditional Independence

- Unconditional (absolute) independence very rare (why?)
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

$$\forall x, y, z : P(x|z, y) = P(x|z)$$

$$X \perp\!\!\!\perp Y | Z$$

- What about this domain:
 - Traffic
 - Umbrella
 - Raining
- What about fire, smoke, alarm?

The Chain Rule Revisited

$$P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$$

The Chain Rule Revisited

$$P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$$

- Trivial decomposition:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain}, \text{Traffic})$$

The Chain Rule Revisited

$$P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$$

- Trivial decomposition:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain}, \text{Traffic})$$

- With assumption of conditional independence:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain})$$

The Chain Rule Revisited

$$P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$$

- Trivial decomposition:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain}, \text{Traffic})$$

- With assumption of conditional independence:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain})$$

- Representation size: 1 + 2 + 4 versus 1 + 2 + 2

The Chain Rule Revisited

$$P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$$

- Trivial decomposition:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain}, \text{Traffic})$$

- With assumption of conditional independence:

$$P(\text{Traffic}, \text{Rain}, \text{Umbrella}) = \\ P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain})$$

- Representation size: 1 + 2 + 4 versus 1 + 2 + 2
- Bayes' nets / graphical models are concerned with distributions with conditional independences